

EXISTENCE OF STATIC VACUUM EXTENSIONS  
FOR NEAR-SCHWARZSCHILD SPHERES: A NEW  
APPROACH TO THE BARTNIK EXTENSION  
PROBLEM

BY

AHMED ELLITHY

A thesis submitted in conformity with  
the requirements for the degree of  
Doctor of Philosophy  
Graduate Department of Mathematics  
University of Toronto

© Copyright 2024 Ahmed Ellithy

# ABSTRACT

---

Existence of Static Vacuum Extensions for Near-Schwarzschild Spheres: A  
New Approach to the Bartnik Extension Problem

Ahmed Ellithy

Doctor of Philosophy

Graduate Department of Mathematics

University of Toronto

2024

We establish the local well-posedness of the Bartnik static metric extension problem for arbitrary Bartnik data that perturb that of any sphere in a Schwarzschild  $\{t = 0\}$  slice. Our result in particular includes spheres with arbitrary small mean curvature. We introduce a new framework to this extension problem by formulating the governing equations in a geodesic gauge, which reduce to a coupled system of elliptic and transport equations. Since standard function spaces for elliptic PDEs are unsuitable for transport equations, we use certain spaces of Bochner-measurable functions traditionally used to study evolution equations. In the process, we establish existence and uniqueness results for elliptic boundary value problems in such spaces in which the elliptic equations are treated as evolutionary equations, and solvability is demonstrated using rigorous energy estimates. The precise nature of the expected difficulty of solving the Bartnik extension problem when the mean curvature is very small is identified and suitably treated in our analysis.

أمي الغالية،  
بفضل تعبك وتضحياتك، أنا وصلت للي أنا فيه. كل حاجة عملتها ليا محفورة في قلبي.

# ACKNOWLEDGEMENTS

---

First and foremost, I would like to express my thanks to my supervisor, Spyros Alexakis. The project you entrusted me with has been a source of a lot of learning and growth. Your patience with my mistakes, your thoughtful feedback on my work, and your support through the most difficult moments of this journey have meant more to me than words can capture.

I am deeply grateful to Professor Yakov Shlapentokh-Rothman for the many enlightening discussions and the invaluable advice and guidance I received. Your kindness and willingness to engage in conversations beyond the academics came at a time when I needed it most. I am fortunate to have had your mentorship.

A special thanks to Almut Burchard, whose compassion and support during my most difficult moments will never be forgotten. You were always there to listen and offer advice when I needed it most, and your empathy made a profound difference in my journey.

To Jemima. You provided an extraordinary amount of support, always ready to listen and offer help. In so many ways, you carried me through the non-academic challenges I faced, and your positivity and encouragement had a huge impact on me.

I am grateful to my friends, whom I will only name a few: Belal, Dev, Jounaid, Nathan, and Doaa. Thank you for always believing in me.

Lastly, I would like to thank my sister for always being there for me, and my mother, without whom I am nothing.

# CONTENTS

---

1	Introduction	1
2	Preliminaries	6
2.1	Properties of Static Vacuum Extensions . . . . .	6
2.2	Function Spaces . . . . .	8
2.3	The Main Theorem . . . . .	12
3	Solvability of Elliptic BVP in $\mathcal{A}_{H_\delta}^{(2,k)}(M)$ and $\mathcal{A}_{C_\delta}^{(2,k)}(M)$	14
4	Reduction of the problem	31
5	Proof of The Main Theorem	37
5.1	Definition of the Artificial Vector Field $X$ . . . . .	38
5.2	Definition and Existence of the Modified Solution . . . . .	43
5.3	The Vanishing of $X$ for Modified solutions $(g, u, X)$ . . . . .	46
5.4	Proof that $D\Phi_{sc}$ is an Isomorphism . . . . .	54
A	Appendix	74
A.1	A PDE of Finite Type for Conformal Killing Vector Fields . . . . .	74
A.2	A Hardy-type Inequality . . . . .	75
A.3	The Legendre functions $P_\ell$ and $Q_\ell$ . . . . .	77

# INTRODUCTION

---

We consider the Bartnik static metric extension problem, which originates in the Bartnik mass-minimization problem, [Bar89, Bar97]: In the latter, one considers a topological 3-ball  $(B, g)$  equipped with a Riemannian 3-metric of positive scalar curvature. A natural example of such a metric arises on any compact space-like maximal hypersurface  $\Omega$  (with boundary) in a  $(3 + 1)$ -spacetime  $(M, g)$  that satisfies the dominant energy condition, where  $g$  is the restriction of the space-time metric  $g$  to  $\Omega$ .

One wishes to assign a notion of mass to  $(\Omega, g)$ ; in fact ideally, [Bar97], the notion of mass should depend just on the restriction of  $g$  to  $\partial\Omega$ , the second fundamental form of  $\Omega \subset M$  at  $\partial\Omega$ , as well as the second fundamental form of  $\partial\Omega$  inside  $\Omega$ .

Bartnik's definition, see [Bar97, Cor17], considers such data on  $\partial\Omega$  and associates to it a class  $\mathcal{PM}$  of admissible asymptotically flat extensions  $(M_{ext}, g_{ext})$ , and seeks to minimize the ADM mass among all such extensions. There are many possibilities on how to define the space  $\mathcal{PM}$  of extensions, see [Bar97]. The most "minimal" requirements are that the 3-metrics  $g_{ext} \in \mathcal{PM}$  should be of positive scalar curvature, the metrics on  $\partial M_{ext}$  induced from the two sides  $\Omega$  (interior side) and  $M_{ext}$  (exterior side) should *match*:  $g_{ext}|_{\partial M_{ext}} = g|_{\partial\Omega}$ ; moreover, the mean curvature  $H_{ext}$  of  $\partial M_{ext}$  in  $M_{ext}$  should agree with the mean curvature  $H_{int}$  of  $\partial M_{ext} = \partial\Omega$  with respect to the interior metric  $g$  over  $\Omega$ . Additional requirements, such as the non-existence of closed minimal surfaces in the extension  $(M_{ext}, g_{ext})$  are very natural (see [Bar97]) and are also frequently imposed. Once the class of admissible extensions has been chosen the Bartnik Mass is defined to be the infimum of the ADM masses, among all admissible extensions.

An important feature of the Bartnik mass is the result of Corvino, [Cor00, Cor17] that if this infimum is attained for some (asymptotically flat) metric  $g$ , on a manifold  $M_{ext}$  with  $\partial M_{ext} = \partial\Omega$ , then this extension  $g$  must satisfy the system of equations:

$$\Delta_g f = 0, \quad \text{Ric}_g = f^{-1} \text{Hess}_g(f), \quad (1.1)$$

as well as the two imposed requirements

$$g|_{\partial M_{ext}} = g|_{\partial\Omega}, \quad H_{ext}(g) = H_{int}. \quad (1.2)$$

A solution to the system (1.1) implies that the metric  $\mathbf{g} = -f^2 dt^2 + \mathfrak{g}$  on  $M_{\text{ext}} \times \mathbb{R}$  would satisfy the Einstein Vacuum equations, and also be static, in the sense that  $\mathcal{L}_{\partial_t} \mathbf{g} = 0$ .

*Remark 1.1.* We note further that if the Bartnik minimizer exists, it is known—see [Barg97]—that the metric  $g_{\text{total}}$  defined over  $M_{\text{total}} = \Omega \cup M_{\text{ext}}$  by joining  $g$  with  $\mathfrak{g}$  across  $\partial M_{\text{ext}}$  is generically expected to be merely Lipschitz across the joining boundary  $\partial M_{\text{ext}}$ : The traceless parts  $\hat{K}|_g, \hat{K}|_{\mathfrak{g}}$  of the second fundamental forms  $K|_g, K|_{\mathfrak{g}}$  induced on  $\partial\Omega = \partial M_{\text{ext}}$  from the two sides  $(\Omega, g), (M_{\text{ext}}, \mathfrak{g})$  are generically expected to not match.

In view of the result of [Coroo] the question of the attainment of the Bartnik mass leads to the Bartnik static extension problem with data supported on a 2-sphere:

**Question** (Bartnik static metric extension problem). *Consider a Riemannian 2-sphere  $(S^2, \gamma)$  equipped with a function  $H$  over  $S^2$ . We consider a (topological) manifold  $M = \mathbb{R}^3 \setminus B$  and seek an asymptotically flat metric  $\mathfrak{g}$  over  $M$  which satisfies:*

- $\mathfrak{g}|_{\partial M} = \gamma$ ,
- the mean curvature  $H_{\text{ext}}$  of  $\partial M$  relative to  $\mathfrak{g}$  equals  $H$ , and:
- There exists a positive function  $f$  over  $M$  with  $f(x) \rightarrow 1$  as  $|x| \rightarrow \infty$  on  $M$  so that the pair  $(\mathfrak{g}, f)$  satisfy the system of equations (1.1).

**Definition 1.** The system in (1.1) will be called the *static vacuum equations*. The pair of prescribed data over  $S^2$  (the metric  $\gamma$  and the putative mean curvature function  $H$ ) will be called *Bartnik data*. A solution  $(\mathfrak{g}, f)$  to (1.1) to this prescribed data with  $\mathfrak{g}$  being asymptotically flat and  $f$  going to 1 at infinity will be called a *static vacuum extension* with Bartnik data  $(\gamma, H)$ .

Two important examples of static vacuum extensions are:

1. The Euclidean solution  $(g_{\text{euc}}, 1)$  on  $\mathbb{R}^3 \setminus B_1$  with Bartnik data  $(\gamma_{S^2}, 1)$ , where  $\gamma_{S^2}$  is the round metric on  $S^2$ .
2. The Riemannian Schwarzschild solution  $(\mathfrak{g}_{\text{sc}}, f_{\text{sc}})$  with mass  $m_0$  on  $\mathbb{R}^3 \setminus B_{r_0}$  and Bartnik data  $(r_0^2 \gamma_{S^2}, \frac{f_{\text{sc}}(r_0)}{r_0})$ , where  $r_0 > 2m_0$  and

$$\mathfrak{g}_{\text{sc}} = f_{\text{sc}}^{-2} dr^2 + r^2 \gamma_{S^2}, \quad f_{\text{sc}} = \sqrt{1 - \frac{2m_0}{r}}$$

Since the static vacuum equations are highly nonlinear, one first hopes to achieve a local well-posedness result near arbitrary solutions. In fact, Anderson and Khuri in [AK09] prove, by means of counterexamples, that global well-posedness does not hold (see also [And23]). Nonetheless, there has been significant progress on establishing local well-posedness results. Miao in [Mia03] confirmed that the extension problem is locally well-posed near Euclidean Bartnik data on the unit sphere under a triple reflectional symmetry assumption. This symmetry assumption was later removed by Anderson in [And15] with the result generalized by Huang and An in [AH21] and [AH22a] for a large range of connected embedded surfaces in Euclidean  $\mathbb{R}^3$ .

Subsequently, Huang and An in [AH22b] introduced a general criterion for local well-posedness near a given solution, which hinges on the triviality of the kernel of a particular operator. They identified a class of static vacuum extensions they call “static regular”, characterized by the linearized operator having a trivial kernel, as sufficient conditions for local well-posedness. They showed that static regularity is, in some sense, generic for smooth hyper surfaces which are already inside a static vacuum extension. However since this relies in a very essential way on the data already lying in the interior of a given solution (which must be analytic), this result does not guarantee genericity in any sense in the space of smooth Bartnik data. Their findings in particular implies that for any given  $m_0 > 0$  and  $\epsilon > 0$ , the set of radii  $r_0 \geq 2m_0 + \epsilon$  for which the Schwarzschild manifold  $\mathbb{R}^3 \setminus B_{r_0}$  with mass  $m_0$  is static regular forms an open dense subset of  $[2m_0 + \epsilon, \infty)$ .

### *Summary of Main Results*

Our main result in this paper is establishing local well-posedness for perturbations of every Schwarzschild solution, hence strengthening Huang and An’s result in [AH22b]. We present a new approach to this problem that can be applied to similar extension problems. In this approach, we write the putative solution  $(g, f)$  with respect to a geodesic gauge, which was not used before for this problem. One benefit of this gauge is that the connection coefficients of the desired solution  $g$  can be linked to  $f$  by *ordinary* differential equations, where  $f$  provides the forcing terms. More precisely, we will be considering the metrics  $g := f^2 \cdot \mathfrak{g}$ , whose Ricci curvature must then satisfy:  $Ric_{ij}(g) = 2\nabla_i u \otimes \nabla_j u$  (with  $u = \ln f$ ), and we will reduce the extension problem in (1.1) to an elliptic equation on  $u$  coupled with Riccati equations on the second fundamental form of  $g$ , with



$u$  providing the forcing term, and constraint equations on the boundary coming from the contracted Gauss and Codazzi equations. We rigorously establish estimates for the linearized operator and its inverse, showing that the linearization of the reduced equations is an isomorphism on appropriate Banach spaces. We then invoke the implicit function theorem on Banach manifolds to conclude local well-posedness.

An interesting remark concerns the expected difficulty of solving Bartnik's extension problem when the Schwarzschild sphere is very close to the horizon, in which the mean curvature is positive but very small. This difficulty is anticipated by the black hole uniqueness theorem (see [Isr67]), which in particular implies the following: for surfaces with zero mean curvature, the existence of static vacuum extensions fails unless the surface is a round sphere, in which case the Schwarzschild exteriors are the only possible extension. Therefore, one expects that the space of allowed perturbations of the Schwarzschild spheres  $S_r$  must be shrinking as  $r$  goes to  $2m_0$ . This also suggests that solving the linearized problem should be progressively harder as  $r \rightarrow 2m_0$ . We do capture this difficulty in our analysis and resolve it (see proposition 5.20), showing the solvability of Bartnik's extension problem near all spheres  $S_r$ ,  $r > 2m_0$ .

The choice of gauge influences which Banach spaces are most appropriate to use. Due to our choice of gauge, the equations in (1.1) reduce to an elliptic PDE coupled with transport equations. Consequently, the standard spaces used for elliptic PDEs, such as weighted Sobolev and Hölder spaces, are not appropriate as they do not provide the correct setting to solve transport equations. Instead, we use spaces of Bochner-measurable functions that are traditionally used as the setting to study hyperbolic and parabolic PDEs (see [Eva98]). More specifically, the spaces we use for  $u$  are  $\mathcal{A}_{C_\delta}^{(2,k)}(M)$  and  $\mathcal{A}_{H_\delta}^{(2,k)}(M)$  defined by (see definition 2.9)

$$u \in \mathcal{A}_{H_\delta}^{(2,k)}(M) \iff \begin{cases} u \in L_\delta^2([r_0, \infty); H^k(S^2)) \\ \partial_r u \in L_{\delta-1}^2([r_0, \infty); H^{k-1}(S^2)) \\ \partial_r^2 u \in L_{\delta-2}^2([r_0, \infty); H^{k-2}(S^2)) \end{cases}$$

$$u \in \mathcal{A}_{C_\delta}^{(2,k)}(M) \iff \begin{cases} u \in C_\delta^0([r_0, \infty); H^k(S^2)) \\ \partial_r u \in C_{\delta-1}^0([r_0, \infty); H^{k-1}(S^2)) \\ \partial_r^2 u \in C_{\delta-2}^0([r_0, \infty); H^{k-2}(S^2)) \end{cases}$$

where  $r_0 > 0$ ,  $k \geq 2$ , and  $\delta \in (-1, -\frac{1}{2})$  is a weight introduced appropriately in the norms of the above spaces to control the decay at infinity. These spaces are not traditionally used to study elliptic PDEs. In this paper, we establish solvability of a certain elliptic problem in the above spaces. More specifically, defining the operator  $Q : u \mapsto (\Delta_g u, u|_{\partial M})$  with respect to a certain asymptotically flat metric  $g$  on  $M = \mathbb{R}^3 \setminus B_{r_0}$ , we demonstrate that (see chapter 3)

$$Q : \mathcal{A}_H^{(2,k)}(M) \rightarrow L_{\delta-2}^2([r_0, \infty); H^{k-2}(S^2)) \times H^{k-1/2}(\partial M) \quad \text{is an isomorphism}$$

$$Q : \mathcal{A}_C^{(2,k)}(M) \rightarrow C_{\delta-2}^0([r_0, \infty); H^{k-2}(S^2)) \times H^k(\partial M) \quad \text{is an isomorphism}$$

This result can be generalized to arbitrary asymptotically flat metrics and more general elliptic boundary value problems, thereby establishing the solvability of such problems in the above spaces.

An interesting comparison we can make to the above is the study of elliptic boundary value problems in  $C^k$  spaces. It is well known that there is no general existence theorem for elliptic boundary value problems in  $C^k(M)$  (see [GT13] problem 4.9 for a counterexample). In [GT13], the authors demonstrate via the celebrated Schauder and Calderon-Zygmund estimates that Hölder spaces  $C^{k,\alpha}(\Omega)$  and Sobolev spaces  $H^k(\Omega)$  on a bounded open set  $\Omega$ , instead of  $C^k(\Omega)$ , have sufficiently nice properties allowing for general existence theorems for elliptic boundary value problems. Modification of those spaces by including weights generalizes these existence results to unbounded spaces (see for example [Bar86]). In this paper, we establish an existence theorem in  $C_{\delta}^0([r_0, \infty); H^k(S^2))$  spaces, a mix of both Hölder and Sobolev spaces. Our work readily implies similar existence results in the spaces  $C^0([a, b] : H^k(S^2))$  when the domain is bounded.

## PRELIMINARIES

---

Let  $M := \mathbb{R}^3 \setminus B_{n \cdot m_0}$  where  $n > 2$  and  $m_0 > 0$ . Denote by  $g_{\text{euc}}$  the Euclidean metric on  $M$  and by  $\gamma_{S^2}$  the round metric on the unit sphere  $S^2$ .

### 2.1 PROPERTIES OF STATIC VACUUM EXTENSIONS

In this section, we will discuss some decay and regularity properties of static vacuum extensions and demonstrate that they can be written in the geodesic gauge. More precisely, we will show that given a static vacuum extension  $(g, f)$ , we can globally write the metric  $g := f^{-2}g$  in geodesic coordinates so that  $g$  takes the form

$$g = f^{-2}dr^2 + \gamma_{g_r}$$

where  $r$  is the distance function from the boundary with respect to  $g$  and  $\gamma_{g_r}$  is the induced metric on the level sets of  $r$ . Note that this form is directly observable in the Schwarzschild solutions  $(g_{\text{sc}}, f_{\text{sc}})$  as  $g_{\text{sc}}$  is given by

$$g_{\text{sc}} = f_{\text{sc}}^{-2}dr^2 + r^2\gamma_{S^2}$$

**Definition 2.1.** Let  $\eta > 0$ . A metric  $g$  over  $M$  is asymptotically flat of order  $\eta > 0$  if there exists a coordinate system  $(x^1, x^2, x^3)$  near infinity in which the metric satisfies

- $g_{ij} - \delta_{ij} = \mathcal{O}(|x|^{-\eta})$
- $\partial_k g_{ij} = \mathcal{O}(|x|^{-\eta-1})$
- $\partial_l \partial_k g_{ij} = \mathcal{O}(|x|^{-\eta-2})$

where  $\partial_k := \frac{\partial}{\partial x^k}$  and  $|x| = \sqrt{|x^1|^2 + |x^2|^2 + |x^3|^2}$ . For conciseness, we will write  $g_{ij} = \delta_{ij} + \mathcal{O}_2(|x|^{-\eta})$  if the above conditions are satisfied.

**Definition 2.2.** For a metric  $g$  and a positive function  $f$ , we say that a pair  $(g, f)$  is a strongly asymptotically flat if  $g$  admits a coordinate system  $(x^1, x^2, x^3)$  near infinity in which

$$\mathfrak{g}_{ij} = \left(1 + \frac{2m}{|x|}\right) \delta_{ij} + \mathcal{O}_2(|x|^{-2}), \quad f = 1 - \frac{m}{|x|} + \mathcal{O}_2(|x|^{-2}) \quad (2.1)$$

The Schwarzschild solutions  $(\mathfrak{g}_{sc}, f_{sc})$  discussed in the introduction are examples of smooth strongly asymptotically flat static vacuum extensions.

The system in (1.1) is equivalent to a lower order system of equations. Letting  $g := f^2 \mathfrak{g}, u := \ln u$ , a direct computation shows that  $(g, f)$  solves equation (1.1) if and only if  $(g, u)$  solve

$$\text{Ric}_g = 2du \otimes du, \quad \Delta_g u = 0. \quad (2.2)$$

We will call the above equations the conformal static vacuum equations. We will call the pair  $(g_{sc}, u_{sc}) := (f_{sc}^2 \mathfrak{g}_{sc}, \ln f_{sc})$  the conformal Schwarzschild solution.

By taking advantage of the form that the Ricci curvature takes for  $g$ , Murchadha in [KM95] shows that every static vacuum extension  $(g, f)$  is strongly asymptotically flat and is smooth away from the boundary. For the rest of this section, we will describe how this strong decay and regularity of static vacuum extensions allows us to write the extension problem in a geodesic gauge.

Let  $(g, u)$  solve the conformal static vacuum equations in (2.2). We wish to write the metric  $g$  in geodesic coordinates. Let  $r(\cdot) = \text{dist}(\cdot, \partial M) + r_0$  be a shifted distance function from  $\partial M$ . Due to the compactness of  $\partial M$ , the function  $r$  is smooth on a neighborhood of  $\partial M$ , with  $\partial M$  excluded, and so defines a foliation near  $\partial M$  with the leaves being the level sets of  $r$ . We can then write the metric with respect to this foliation as  $dr^2 + \gamma_r$  where  $\gamma_r$  is the induced metric on the level sets. This representation of the metric generally does not hold globally and is valid only whenever  $r$  is differentiable. However, under a smallness assumption on the Ricci curvature of  $g$ , it will hold that  $r$  is differentiable everywhere on  $M \setminus \partial M$  and the metric can be written globally as  $dr^2 + \gamma_r$ . This follows from the next proposition, which follows from a straightforward adaptation of the argument in proposition 5.01 in [CK93].

**Proposition 2.3.** *There exists  $\tau' = \tau'(n, m_0) > 0$  small enough such that the following is true for any  $0 < \tau < \tau'$ .*

*If an asymptotically flat metric  $g$  on  $M$  of order  $\eta > 0$  satisfies  $|\text{Ric}_g| < \tau \text{dist}_g(x, \partial M)^{-2-\eta}$  for  $x \in M$ , then:*

1. The affine parameter  $r(\cdot) = \text{dist}_g(\partial M, \cdot) + r_0$  is differentiable everywhere on  $M \setminus \partial M$  and defines a global radial foliation with leaves  $S_r$  diffeomorphic to  $S^2$ . Moreover, given a coordinate system  $(x^1, x^2, x^3)$  near infinity as described in definition 2.1,  $r$  and  $|x|$  are comparable in the sense that

$$C^{-1}|x| \leq r \leq C|x| \quad (2.3)$$

for some constant  $C > 0$ .

2. With respect to this foliation, we have

$$\text{tr}K = \frac{2}{r} + \mathcal{O}_1(r^{-1-\eta}), \quad |\hat{K}| = \mathcal{O}_1(r^{-1-\eta}) \quad (2.4)$$

where  $K = \text{Hess}(r)$  is the second fundamental form on the leaves  $S_r$ ,  $\text{tr}K$  is the trace of  $K$ , and  $\hat{K}$  is the traceless part of  $K$ .

3. There exists a unique diffeomorphism  $\Phi : M \rightarrow [nm_0, \infty) \times S^2$  such that  $\Phi|_{\partial M} = \text{Id}_{S^2}$ ,  $r(\cdot) = \pi_r \circ \Phi(\cdot)$  where  $\pi_r$  is the projection onto the first coordinate, and  $\Phi_*g = dr^2 + \gamma_{g_r}$  where  $\gamma_{g_r}$  is the push forward of the induced metric on  $S_r$ .

This allows us to globally express the static vacuum extension  $(g, f)$  in geodesic coordinates as follows:

$$g = f^{-2}dr^2 + \gamma_{g_r}, \quad \text{where } \gamma_{g_r} \text{ is the induced metric on } S_r$$

and, hence, justifies the space of metrics that we will be working in (see definition 2.8 in the next section).

## 2.2 FUNCTION SPACES

In this section, we define the function spaces that we will be using. Fix  $k \in \mathbb{Z}_{\geq 0}$  and  $\delta \in \mathbb{R}$ . From here onwards, we will identify  $M$  with the space  $[nm_0, \infty) \times S^2$ .

**Definition 2.4.** We define the weighted Sobolev space  $H_\delta^k(M)$  with weight  $\delta$  to be the space of all functions  $u$  in  $H_{loc}^k(M)$  such that  $\|u\|_{k,\delta} < \infty$  respectively, where

$$\|u\|_{k,\delta} = \sum_{l=0}^k \left\{ \int_M (|D^l u| \cdot r^{l-\delta})^2 r^{-3} dV \right\}^{\frac{1}{2}} \quad (2.5)$$

where  $r = |x|$ ,  $D$  is the connection with respect to the Euclidean metric on  $M$ , and  $dV$  is the Euclidean volume form on  $M$ . We will also denote the

space  $H_\delta^k(M)$  by  $L_\delta^2(M)$  when  $k = 0$ .

**Definition 2.5.** We define the space  $\mathcal{X}_\delta^k(M)$  to be the space of vector fields  $X$  on  $M$  with components  $X^i := X(x^i)$  in  $H_\delta^k(M)$ , where  $(x^1, x^2, x^3)$  is the standard cartesian coordinates. The norm we use is

$$\|X\|_{k,\delta} := \sum_{l=0}^k \left\| |D^l X| \right\|_{0,\delta-l} \quad (2.6)$$

**Definition 2.6.** Let  $H^k(S^2)$  be the usual  $L^2$  space, when  $k = 0$ , and Sobolov space, when  $k \geq 1$ , on  $(S^2, \gamma_{S^2})$ . Let  $\mathcal{M}^k(S^2)$  and  $\mathcal{H}^k(S^2)$  be the space of metrics on  $S^2$  and symmetric tensors on  $S^2$ , respectively, with components in  $H^k(S^2)$ . The norm we will use is as follows:

$$\|h\|_{\mathcal{H}^k(S^2)}^2 := \sum_{l=0}^k \left\| |D^l h| \right\|_{L^2(S^2)}^2 \quad (2.7)$$

where  $D$  is the covariant derivative on  $S^2$  with respect  $\gamma_{S^2}$ .

Let  $\Omega^k(S^2)$  be the space of  $k$ -forms on  $S^2$  with components in  $H^k(S^2)$ . The norm used on this space is as follows:

$$\|\omega\|_{\Omega^k(S^2)}^2 := \sum_{l=0}^k \left\| |D^l \omega| \right\|_{L^2(S^2)}^2 \quad (2.8)$$

**Definition 2.7.** Let  $t \in \mathbb{Z}_{\geq 0}$ . We define the space  $H_\delta^t([nm_0, \infty); H^k(S^2))$  to be the space of functions  $u$  in  $H_{loc}^t([nm_0, \infty); H^k(S^2))$  such that  $\|u\|_{H,(t \rightarrow k),\delta} < \infty$ , where

$$\|u\|_{H,(t \rightarrow k),\delta}^2 := \sum_{t'=0}^t \int_{nm_0}^{\infty} r^{-2\delta-1+2t'} \left\| \partial_r^{(t')} u(r) \right\|_{H^k(S^2)}^2 dr \quad (2.9)$$

We also define the space  $C_\delta^t([nm_0, \infty); H^k(S^2))$  to be the space of continuous  $H^k(S^2)$ -valued functions  $u$  on  $[nm_0, \infty)$  such that  $\|u\|_{C,(t \rightarrow k),\delta} < \infty$ , where

$$\|u\|_{C,(t \rightarrow k),\delta}^2 := \sum_{t'=0}^t \sup_{r \geq nm_0} \left( r^{-2\delta+2t'} \left\| \partial_r^{(t')} u(r) \right\|_{H^k(S^2)}^2 \right) \quad (2.10)$$

We then define the space  $H_\delta^t([nm_0, \infty); \mathcal{M}^k(S^2))$  and  $H_\delta^t([nm_0, \infty); \mathcal{H}^k(S^2))$  similarly to the above with norm

$$\|h\|_{H_\delta^t([nm_0, \infty); \mathcal{H}^k(S^2))}^2 := \sum_{t'=0}^t \int_{nm_0}^\infty r^{-2\delta-1+2t'} \left\| \partial_r^{(t')} h(r) \right\|_{\mathcal{H}^k(S^2)}^2 dr \quad (2.11)$$

**Definition 2.8.** Define  $\mathcal{M}_\delta^k(M)$  to be the space of metrics on  $[nm_0, \infty) \times S^2$  of the form  $dr^2 + g(r)$  where  $g(r) = r^2(\gamma_\infty + h(r))$ ,  $\gamma_\infty \in \mathcal{M}^k(S^2)$ , and  $h \in H_\delta^2([nm_0, \infty); \mathcal{H}^k(S^2))$ . The space  $\mathcal{M}_\delta^k(M)$  can be naturally identified with an open subset of the Banach space  $\mathcal{H}^k(S^2) \oplus H_\delta^2([nm_0, \infty); \mathcal{H}^k(S^2))$ . This makes  $\mathcal{M}_\delta^k(M)$  an open Banach submanifold of  $\mathcal{H}^k(S^2) \oplus H_\delta^2([nm_0, \infty); \mathcal{H}^k(S^2))$  and, in particular, a Banach manifold. Given  $g_0 \in \mathcal{M}_\delta^k(M)$ , the tangent space  $T_{g_0} \mathcal{M}_\delta^k(M)$  is isomorphic to the space of tensors  $\tilde{g}$  of the form  $\tilde{g} = r^2(\tilde{\gamma}_\infty + \tilde{h}(r))$ , where  $\tilde{\gamma}_\infty \in \mathcal{H}^k(S^2)$  and  $\tilde{h} \in H_\delta^2([nm_0, \infty); \mathcal{H}^k(S^2))$ , equipped with the norm

$$\|\tilde{g}\|_{\mathcal{M}_\delta^k} := \|\tilde{\gamma}_\infty\|_{\mathcal{H}^k(S^2)} + \|\tilde{h}\|_{H_\delta^2([nm_0, \infty); \mathcal{H}^k(S^2))} \quad (2.12)$$

**Definition 2.9.** Let  $t \geq 0$ . Denote by  $\mathcal{A}_{H_\delta}^{(t,k)}(M)$  and  $\mathcal{A}_{C_\delta}^{(t,k)}(M)$  the spaces

$$\mathcal{A}_{H_\delta}^{(t,k)}(M) := \bigcap_{t'=0}^t H_\delta^{t'}([nm_0, \infty); H^{k-t'}(S^2)), \quad \mathcal{A}_{C_\delta}^{(t,k)}(M) := \bigcap_{t'=0}^t C_\delta^{t'}([nm_0, \infty); H^{k-t'}(S^2)) \quad (2.13)$$

equipped with the norms

$$\|u\|_{\mathcal{A}_{H_\delta}^{(t,k)}}^2 := \max_{0 \leq t' \leq t} \|u\|_{H_\delta^{t'}([nm_0, \infty); H^{k-t'}(S^2))}^2, \quad \|u\|_{\mathcal{A}_{C_\delta}^{(t,k)}}^2 := \max_{0 \leq t' \leq t} \|u\|_{C_\delta^{t'}([nm_0, \infty); H^{k-t'}(S^2))}^2 \quad (2.14)$$

Note that

$$u \in \mathcal{A}_{H_\delta}^{(t,k)}(M) \iff \text{for every } 0 \leq t' \leq t, \partial_r^{(t')} u \in L_{\delta-t'}^2([nm_0, \infty); H^{k-t'}(S^2))$$

$$u \in \mathcal{A}_{C_\delta}^{(t,k)}(M) \iff \text{for every } 0 \leq t' \leq t, \partial_r^{(t')} u \in C_{\delta-t'}^0([nm_0, \infty); H^{k-t'}(S^2))$$

Denote the intersection of these spaces by  $\mathcal{A}_\delta^{(t,k)}(M)$  defined by

$$\mathcal{A}_\delta^{(t,k)}(M) := \mathcal{A}_{H_\delta}^{(t,k)}(M) \cap \mathcal{A}_{C_\delta}^{(t,k)}(M)$$

equipped with the norm

$$\|u\|_{\mathcal{A}_\delta^{(t,k)}}^2 := \max_{0 \leq t' \leq t} (\|u\|_{H, (t' \rightarrow k - t'), \delta}^2 + \|u\|_{C, (t' \rightarrow k - t'), \delta}^2) \quad (2.15)$$

In the next proposition, we list some important results regarding the spaces we defined that will be repeatedly used in the rest of the paper.

**Proposition 2.10.**

- (a) Let  $k \geq 0$ ,  $t \geq 1$  and  $\delta < 0$ . Every function  $u \in H_\delta^t([nm_0, \infty); H^k(S^2))$  has a representative in  $C_{loc}^{t-1}([nm_0, \infty); H^k(S^2))$ , which will also be denoted by  $u$ . Furthermore, there exists a constant  $C > 0$  such that for every  $u \in H_\delta^t([nm_0, \infty); H^k(S^2))$

$$\|u\|_{C, (t-1 \rightarrow k), \delta} \leq C \|u\|_{H, (t \rightarrow k), \delta} \quad (2.16)$$

If in addition  $k \geq 2$ , then  $\partial_r^{(t')} u(r) \in C^{k-2}(S^2)$  for every  $r \in [nm_0, \infty)$  and  $0 \leq t' \leq t-1$ . Also, for every  $0 \leq l \leq k-2$  and  $0 \leq t' \leq t-1$ ,

$$|\mathcal{D}^l \partial_r^{(t')} u| = o(r^{\delta-l-t'}) \quad \text{as } r \rightarrow \infty \quad (2.17)$$

- (b) Let  $dr^2 + g(r) \in \mathcal{M}_\delta^k(M)$  with  $g(r) = r^2(\gamma_\infty + h(r))$ .

Then with respect to the foliation defined by the level sets of  $r$ , the trace and traceless part of the fundamental form satisfy

$$\left| \text{tr}K - \frac{2}{r} \right| \in H_{\delta-1}^1([nm_0, \infty); H^k(S^2)), \quad \hat{K} \in H_{\delta-1}^1([nm_0, \infty); \mathcal{H}^k(S^2)) \quad (2.18)$$

Furthermore, the metric  $dr^2 + g(r)$  is asymptotically flat if and only if  $\gamma_\infty$  is of constant curvature 1.

- (c) Let  $k_1 < k_2$ ,  $t_1 < t_2$ , and  $\delta_2 < \delta_1 < 0$ . Then the space  $H_{\delta_2}^{t_2}([nm_0, \infty); H^{k_2}(S^2))$  is compactly embedded in  $H_{\delta_1}^{t_1}([nm_0, \infty); H^{k_1}(S^2))$ . Furthermore, the space  $C_{\delta_2}^{t_2}([nm_0, \infty); H^{k_2}(S^2))$  is compactly embedded in  $C_{\delta_1}^{t_1}([nm_0, \infty); H^{k_1}(S^2))$ .
- (d) Let  $k \geq 1$  and  $\delta \in \mathbb{R}$ . Suppose a function  $u$  satisfies

$$u \in L_\delta^2([nm_0, \infty); H^k(S^2))(M), \quad \partial_r u \in L_{\delta-1}^2([nm_0, \infty); H^{k-1}(S^2))(M)$$

Then for every  $r \in [nm_0, \infty)$ ,

$$u(r) \in H^{k-1/2}(S^2)$$



*Proof.* (a), (b) and (d) follow immediately from standard results on Sobolev spaces (see [Bar86] theorem 1.2 and lemma 1.4, and [Eva98] section 5.9).

We focus on proving (b). The metric  $g(r)$  evolves according to the equation

$$\partial_r g(r) = \text{tr}K g(r) + 2\hat{K} \quad (2.19)$$

where  $K = \text{Hess}(r)$  is the second fundamental form on the leaves  $S_r$ ,  $\text{tr}K$  is the trace of  $K$ , and  $\hat{K}$  is the traceless part of  $K$ . Since  $g(r) = r^2(\gamma_\infty + h(r))$ , it follows that

$$\text{tr}K = \frac{2}{r} + \frac{1}{2}\text{tr}_{r^{-2}g(r)}(\partial_r h(r)), \quad 2\hat{K} = r^2\left(\partial_r h(r) - \text{tr}_{g(r)}(\partial_r h(r))g(r)\right) \quad (2.20)$$

This directly implies equation (2.18)

In view of the Gauss and Codazzi equations, we get

$$R = 2\text{Ric}\left(\frac{\partial}{\partial r}, \frac{\partial}{\partial r}\right) + R_{S_r} - \frac{1}{2}\text{tr}K^2 + |\hat{K}|^2, \quad \text{Ric}\left(\frac{\partial}{\partial r}, \frac{\partial}{\partial r}\right) = -\partial_r \text{tr}K - \frac{1}{2}(\text{tr}K)^2 - |\hat{K}|^2 \quad (2.21)$$

It follows immediately using equation (2.18) and the fact that the scalar curvature  $R_{\gamma_\infty}$  of  $(S^2, \gamma_\infty)$  is the limit of  $r^2 R_{S_r}$  as  $r$  goes to infinity that  $R$  decays faster than  $r^{-2}$  if and only if  $R_{\gamma_\infty} = 2$ . We conclude that the metric is asymptotically flat if and only if  $\gamma_\infty$  is of constant curvature 1.  $\square$

### 2.3 THE MAIN THEOREM

**Definition 2.11.** In place of the mean curvature  $H$ , we will work with  $\text{tr}K_{\mathfrak{B}} := 2H$  for convenience, which represents the trace of the hypothetical second fundamental form on  $\partial M$ . From this point forward, we will denote the Bartnik data on  $\partial M$  by  $(\gamma_{\mathfrak{B}}, \frac{1}{2}\text{tr}K_{\mathfrak{B}})$ . We will also denote by  $(\gamma_{\mathfrak{g}_{\text{sc}}}, \frac{1}{2}\text{tr}K_{\mathfrak{g}_{\text{sc}}})$  the Schwarzschild Bartnik data, which is given by

$$\gamma_{\mathfrak{g}_{\text{sc}}} = (nm_0)^2 \gamma_{S^2}, \quad \text{tr}K_{\mathfrak{g}_{\text{sc}}} = \frac{2\sqrt{1 - \frac{2}{n}}}{nm_0} \quad (2.22)$$

The statement of the main theorem is as follows.

**Main Theorem.** *Let  $\delta \in (-1, -\frac{1}{2}]$  and  $k \geq 5$ . There exists a neighbourhood  $\mathcal{U}$  of  $(\gamma_{\mathfrak{g}_{\text{sc}}}, \frac{1}{2}\text{tr}K_{\mathfrak{g}_{\text{sc}}})$  in  $\mathcal{M}^{k+1}(\partial M) \times H^k(\partial M)$  and a unique  $\mathcal{C}^1$  map*

$\mathbf{H} : (\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}}) \mapsto (g, u)$  on  $\mathcal{U}$  to  $\mathcal{M}_{\delta}^k(M) \times \mathcal{A}_{\delta}^{(2,k+1)}(M)$  in which  $(\mathfrak{g}, f) := (e^{-2u}g, e^u)$  solves the static Einstein vacuum equations with Bartnik data  $(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}})$ .

Given Bartnik data  $(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}}) \in \mathcal{U}$ , the pair  $(g, u) = \mathbf{H}(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}})$  will then solve the conformal static vacuum equations written out in equation (2.2). Due to proposition 2.10,  $g$  and  $f$  are  $\mathcal{C}^2$  on  $M$  and satisfy, in some coordinates  $(x^1, x^2, x^3)$ ,

$$\mathfrak{g}_{ij} = \delta_{ij} + \mathcal{O}_2(|x|^{\delta}), \quad f = 1 + \mathcal{O}_2(|x|^{\delta}) \quad (2.23)$$

Moreover, the discussion in section 2.1 implies that  $(\mathfrak{g}, f)$  is strongly asymptotically flat and is smooth away from the boundary.

# SOLVABILITY OF ELLIPTIC BVP IN $\mathcal{A}_{H_\delta}^{(2,k)}(M)$ AND $\mathcal{A}_{C_\delta}^{(2,k)}(M)$

---

In this chapter, we will establish the well-posedness of the elliptic PDE  $\Delta_{g_{sc}} \tilde{u} = 0$  on  $(M, g_{sc})$ , subject to Dirichlet boundary conditions, in the function spaces  $\mathcal{A}_{H_\delta}^{(2,k)}(M)$  and  $\mathcal{A}_{C_\delta}^{(2,k)}(M)$ . Here,  $g_{sc}$  is the conformal Schwarzschild metric on  $M = \mathbb{R}^3 \setminus B_{n \cdot m_0}$  given by

$$g_{sc} = dr^2 + r(r - 2m_0)\gamma_{S^2} \quad (3.1)$$

and  $n > 2$ ,  $m_0 > 0$ .

More precisely, we will prove the following theorem.

**Theorem 3.1.** *Define the operator  $\mathcal{Q}$  by:*

$$\mathcal{Q}(\tilde{u}) := \begin{pmatrix} \Delta_{g_{sc}} \tilde{u} \\ \tilde{u} \end{pmatrix}$$

For  $\delta \in (-1, -\frac{1}{2}]$  and  $k \geq 1$ ,

$$\mathcal{Q} : \mathcal{A}_{H_\delta}^{(2,k+1)}(M) \rightarrow \mathcal{A}_{H_{\delta-2}}^{(0,k-1)}(M) \times H^{k+1/2}(\partial M) \quad \text{is an isomorphism}$$

and

$$\mathcal{Q} : \mathcal{A}_{C_\delta}^{(2,k+1)}(M) \rightarrow \mathcal{A}_{C_{\delta-2}}^{(0,k-1)}(M) \times H^{k+1}(\partial M) \quad \text{is an isomorphism}$$

*Remark 3.2.* In particular, it holds that

$$\mathcal{Q} : \mathcal{A}_\delta^{(2,k+1)}(M) \rightarrow \mathcal{A}_{\delta-2}^{(0,k-1)}(M) \times H^{k+1}(\partial M) \quad \text{is an isomorphism,}$$

which will be used in section 5.4.

The map  $\mathcal{Q}$  can be defined on the space  $\mathcal{A}_{H_\delta}^{(2,k+1)}(M)$  and  $\mathcal{A}_{C_\delta}^{(2,k+1)}(M)$  with codomain  $\mathcal{A}_{H_{\delta-2}}^{(0,k-1)}(M) \times H^{k+1/2}(\partial M)$  and  $\mathcal{A}_{C_{\delta-2}}^{(0,k-1)}(M) \times H^{k+1}(\partial M)$

respectively. Indeed, we deduce directly from the definition of our Banach spaces that for all  $u \in \mathcal{A}_{H_\delta}^{(2,k+1)}(M)$ ,

$$\Delta_{g_{sc}} \tilde{u} = \partial_r^2 \tilde{u} + \frac{2(r-m_0)}{r(r-2m_0)} \partial_r \tilde{u} + \frac{1}{r(r-2m_0)} \mathbb{A}_{\gamma_{S^2}} \tilde{u}(r) \in L_{\delta-2}^2 \left( [nm_0, \infty); H^{k-1}(S^2) \right) = \mathcal{A}_{H_{\delta-2}}^{(0,k-1)}(M)$$

$$\tilde{u}(nm_0) \in H^{k+1/2}(S^2) \quad (\text{by proposition 2.10 (d)})$$

Similarly, for all  $u \in \mathcal{A}_{C_\delta}^{(2,k+1)}(M)$ ,

$$\Delta_{g_{sc}} \tilde{u} = \partial_r^2 \tilde{u} + \frac{2(r-m_0)}{r(r-2m_0)} \partial_r \tilde{u} + \frac{1}{r(r-2m_0)} \mathbb{A}_{\gamma_{S^2}} \tilde{u}(r) \in C_{\delta-2}^0 \left( [nm_0, \infty); H^{k-1}(S^2) \right) = \mathcal{A}_{C_{\delta-2}}^{(0,k-1)}(M)$$

$$\tilde{u}(nm_0) \in H^{k+1}(S^2)$$

We recall the following result from Maxwell in [Max05]:

$$Q : H_\delta^2(M) \rightarrow L_{\delta-2}^2(M) \times H^{3/2}(\partial M) \quad \text{is an isomorphism}$$

To prove theorem 3.1, it then suffices to prove the estimates

$$\|\tilde{u}\|_{\mathcal{A}_{H_\delta}^{(2,k+1)}} \leq C \|Q(\tilde{u})\|_{\mathcal{A}_{H_{\delta-1}}^{(0,k-1)} \times H^{k+1/2}(\partial M)}$$

$$\|\tilde{u}\|_{\mathcal{A}_{C_\delta}^{(2,k+1)}} \leq C \|Q(\tilde{u})\|_{\mathcal{A}_{C_{\delta-1}}^{(0,k-1)} \times H^{k+1}(\partial M)}$$

for all  $\tilde{u}$  in  $\mathcal{A}_{H_\delta}^{(2,k+1)}(M)$  and  $\mathcal{A}_{C_\delta}^{(2,k+1)}(M)$  respectively. These estimates will be the content of the next lemma.

**Lemma 3.3.** • *There exist a constant  $C > 0$  such that for any  $\tilde{u} \in \mathcal{A}_{H_\delta}^{(2,k+1)}(M)$ , the following estimate holds.*

$$\|\tilde{u}\|_{\mathcal{A}_{H_\delta}^{(2,k+1)}} \leq C \left( \|\Delta_{g_{sc}} \tilde{u}\|_{\mathcal{A}_{H_\delta}^{(0,k-1)}} + \|\tilde{u}\|_{H^{k+1/2}(\partial M)} \right) \quad (3.2)$$

• *There exist a constant  $C > 0$  such that for any  $\tilde{u} \in \mathcal{A}_{C_\delta}^{(2,k+1)}(M)$ , the following estimate holds.*

$$\|\tilde{u}\|_{\mathcal{A}_{C_\delta}^{(2,k+1)}} \leq C \left( \|\Delta_{g_{sc}} \tilde{u}\|_{\mathcal{A}_{C_\delta}^{(0,k-1)}} + \|\tilde{u}\|_{H^{k+1}(\partial M)} \right) \quad (3.3)$$

*Proof.* Since  $\mathcal{A}_\delta^{(2,k+1)}(M)$  is dense in both  $\mathcal{A}_{H_\delta}^{(2,k+1)}(M)$  and  $\mathcal{A}_{C_\delta}^{(2,k+1)}(M)$ , it suffices to prove both estimates for all  $\tilde{u} \in \mathcal{A}_\delta^{(2,k+1)}(M)$ .

Let  $\tilde{u} \in \mathcal{A}_\delta^{(2,k+1)}$ . Define  $F := \Delta_{g_{sc}} \tilde{u}$  and  $h := \tilde{u}(nm_0)$ . Then

$$F \in \mathcal{A}_{\delta-2}^{(0,k-1)}(M) = L_{\delta-2}^2\left([nm_0, \infty); H^{k-1}(S^2)\right) \cap C_{\delta-2}^0\left([nm_0, \infty); H^{k-1}(S^2)\right), \quad h \in H^{k+1}(\partial M)$$

We utilize the spherical symmetry of  $(M, g_{sc})$  to reduce the equation  $\Delta_{g_{sc}} \tilde{u} = F$  to differential equations on the coefficients of  $\tilde{u}$  with respect to its spherical harmonics decomposition. Decompose  $\tilde{u}$ ,  $F$ , and  $h$  as follows

$$\tilde{u}(r, x) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} a_{m\ell}(r) Y_{m\ell}(x), \quad F(r, x) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} b_{m\ell}(r) Y_{m\ell}(x), \quad h(x) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} c_{m\ell} Y_{m\ell}(x) \quad (3.4)$$

for  $r \in [nm_0, \infty)$  and  $x \in \partial M$ . Here and below the spherical harmonics  $Y_{m\ell}(x)$  are viewed as functions over the unit sphere  $S^2$ . These same functions will also be thought of over round spheres of any other radius via the natural push-forward map. We will assume that they are normalized with respect to the round metric  $\gamma_{S^2}$  on the unit sphere.

We first rewrite the norms of the relevant Banach spaces in terms of the coefficients with respect to the spherical harmonics decomposition. For a nonnegative integer  $s$  and  $f \in H^s(\partial M)$  with spherical harmonic coefficients  $f_{m\ell}$ , the norm

$$\|f\|_{H^s(\partial M)} := \left( \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^s |f_{m\ell}|^2 \right)^{1/2} \quad (3.5)$$

is equivalent to the standard norm on  $H^s(\partial M)$ . For a nonnegative integer  $t$  and real number  $\tau$ , we will rewrite the norm on  $H_\tau^t([nm_0, \infty); H^s(S^2))(M)$  and  $C_\tau^t([nm_0, \infty); H^s(S^2))$ .

We begin with the norm on  $H_\tau^t([nm_0, \infty); H^s(S^2))(M)$ . Given a function  $v \in H_\tau^t([nm_0, \infty); H^s(S^2))$ , recall that

$$\|v\|_{H_{(t \rightarrow s), \tau}^t}^2 = \sum_{t'=0}^t \int_{nm_0}^{\infty} r^{-2\delta-1+2t'} \left\| \partial_r^{t'} v(r) \right\|_{H^s(S^2)}^2 dr \quad (3.6)$$

Let  $v_{m\ell} = v_{m\ell}(r)$  be the spherical harmonic coefficients for  $v$ . The first term in the sum becomes

$$\int_{nm_0}^{\infty} r^{-2\delta-1} \|v(r)\|_{H^s(S^2)}^2 dr = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell+1)]^s \int_{nm_0}^{\infty} r^{-2\delta-1} |v_{m\ell}(r)|^2 dr \quad (3.7)$$

where we invoked the monotone convergence theorem to switch the order of the integral and the infinite sum. Now for each  $1 \leq t' \leq t$  and  $r \in [nm_0, \infty)$ , we have that

$$\int_{S^2} Y_{m\ell} \partial_r^{t'} v(r) d\sigma_{S^2} = \partial_r^{t'} \int_{S^2} Y_{m\ell} v(r) d\sigma_{S^2} \quad (3.8)$$

since  $\partial_r^{t'-1}$  lives in  $H_{loc}^1([nm_0, \infty); L^2(S^2))$ . In light of the fact that  $v_{m\ell}(r) = \int_{S^2} Y_{m\ell} v(r) d\sigma_{S^2}$ , it follows that  $v_{m\ell}$  are differentiable  $t$  times in  $r$  and  $v_{m\ell}^{(t')}$  are the spherical harmonic coefficients of  $\partial_r^{t'} v$ .

We can then rewrite the norm in equation (3.6) as follows:

$$\|v\|_{(t \rightarrow s), \tau}^2 = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell+1)]^s \sum_{t'=0}^t \int_{nm_0}^{\infty} r^{-2\tau-1+2t'} \left(v_{m\ell}^{(t')}(r)\right)^2 dr \quad (3.9)$$

where we have repeatedly invoked the monotone convergence theorem to switch the order of the integral and the infinite sum.

Now consider a function  $w \in C_{\tau}^t([nm_0, \infty); H^s(S^2))$ . Recall that

$$\|w\|_{C, (t \rightarrow s), \tau}^2 = \sum_{t'=0}^t \sup \left( r^{-2\tau+2t'} \left\| \partial_r^{t'} w(r) \right\|_{H^s(S^2)}^2 \right) \quad (3.10)$$

Letting  $w_{m\ell} = w_{m\ell}(r)$  be the spherical harmonic coefficients for  $w$ , we have

$$\|w\|_{C, (t \rightarrow s), \tau}^2 = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell+1)]^s \sum_{t'=0}^t \sup \left( r^{-2\tau+2t'} (w_{m\ell}^{(t')}(r))^2 \right) \quad (3.11)$$

It is then convenient to define the following norms for functions on  $[nm_0, \infty)$ : for a function  $f_1 \in H_{loc}^t([nm_0, \infty))$  and  $f_2 \in C^t([nm_0, \infty))$ , define

$$\|f_1\|_{H, t, \tau}^2 := \sum_{t'=0}^t \int_{nm_0}^{\infty} r^{-2\tau-1+2t'} \left(f_1^{(t')}(r)\right)^2 dr, \quad \|f_2\|_{C, t, \tau}^2 := \sum_{t'=0}^t \sup \left( r^{-2\tau+2t'} (f_2^{(t')}(r))^2 \right) \quad (3.12)$$

We denote by  $H_\tau^t([nm_0, \infty))$  and  $C_\tau^t([nm_0, \infty))$  all functions  $f_1 \in H_{loc}^t([nm_0, \infty))$  and  $f_2 \in C^t([nm_0, \infty))$  in which  $\|f_1\|_{H,t,\tau} < \infty$  and  $\|f_2\|_{C,t,\tau} < \infty$  respectively.

Using the above notation, we can then write equation (3.6) and (3.11) as follows:

$$\begin{aligned} \|v\|_{H,(t \rightarrow s),\tau}^2 &= \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^s \|v_{m\ell}\|_{H,t,\tau}^2 \\ \|w\|_{C,(t \rightarrow s),\tau}^2 &= \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^s \|w_{m\ell}\|_{C,t,\tau}^2 \end{aligned} \quad (3.13)$$

We now return to the statement of the lemma. Recall from equation (3.4) that

- $\tilde{u} \in \mathcal{A}_\delta^{(2,k+1)}(M)$  with coefficients  $a_{m\ell} \in H_\delta^2([nm_0, \infty)) \cap C_\delta^2([nm_0, \infty))$ .
- $F := \Delta_{g_{sc}} \tilde{u} \in \mathcal{A}_{\delta-2}^{(0,k-1)}(M)$  with coefficients  $b_{m\ell} \in L_{\delta-2}^2([nm_0, \infty)) \cap C_{\delta-2}([nm_0, \infty))$ .
- $h := \tilde{u}(nm_0) \in H^{k+1}(S^2)$  with coefficients  $c_{m\ell}$ .

To prove the lemma, it then suffices to show that there exist a constant  $C > 0$  independent of  $m$  and  $\ell$  such that

$$\begin{aligned} \|a''_{m\ell}\|_{H,0,\delta-2}^2 + [1 + \ell(\ell + 1)] \|a'_{m\ell}\|_{H,0,\delta-1}^2 + [1 + \ell(\ell + 1)]^2 \|a_{m\ell}\|_{H,0,\delta}^2 \\ \leq C(\|b_{m\ell}\|_{H,0,\delta-2}^2 + [1 + \ell(\ell + 1)]^{3/2} |c_{m\ell}|^2) \end{aligned} \quad (\mathbf{H-Est})$$

$$\begin{aligned} \|a''_{m\ell}\|_{C,0,\delta-2}^2 + [1 + \ell(\ell + 1)] \|a'_{m\ell}\|_{C,0,\delta-1}^2 + [1 + \ell(\ell + 1)]^2 \|a_{m\ell}\|_{C,0,\delta}^2 \\ \leq C(\|b_{m\ell}\|_{C,0,\delta-2}^2 + [1 + \ell(\ell + 1)]^2 |c_{m\ell}|^2) \end{aligned} \quad (\mathbf{C-Est})$$

for each  $m$  and  $\ell$ . Indeed, if we multiply **H-Est** and **C-Est** by  $[1 + \ell(\ell + 1)]^{k-1}$  and sum over  $m$  and  $\ell$ , we get the two desired estimates in the statement of the lemma. We will demonstrate this for **H-Est**: after

multiplying **H-Est** by  $[1 + \ell(\ell + 1)]^{k-1}$ , we sum over  $m$  and  $\ell$  to get the following three estimates

$$\begin{aligned}
 & \|\tilde{u}\|_{H,(2 \rightarrow k-1),\delta}^2 \\
 &= \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^{k-1} \|a_{m\ell}\|_{H,2,\delta}^2 \\
 &= \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^{k-1} \left( \|a''_{m\ell}\|_{H,0,\delta-2}^2 + \|a'_{m\ell}\|_{H,0,\delta-1}^2 + \|a_{m\ell}\|_{H,0,\delta}^2 \right) \\
 &\leq C \left( \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^{k-1} \|b_{m\ell}\|_{H,0,\delta-2}^2 + \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^{k+1/2} |c_{m\ell}|^2 \right) \\
 &= C \left( \|F\|_{\mathcal{A}_{H\delta-2}^{(0,k-1)}}^2 + \|h\|_{H^{k+1/2}(S^2)}^2 \right)
 \end{aligned}$$

$$\begin{aligned}
 & \|\tilde{u}\|_{H,(1 \rightarrow k),\delta}^2 \\
 &= \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^k \|a_{m\ell}\|_{H,1,\delta}^2 \\
 &= \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^k \left( \|a'_{m\ell}\|_{H,0,\delta-1}^2 + \|a_{m\ell}\|_{H,0,\delta}^2 \right) \\
 &\leq C \left( \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^{k-1} \|b_{m\ell}\|_{H,0,\delta-2}^2 + \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^{k+1/2} |c_{m\ell}|^2 \right) \\
 &= C \left( \|F\|_{\mathcal{A}_{H\delta-2}^{(0,k-1)}}^2 + \|h\|_{H^{k+1/2}(S^2)}^2 \right)
 \end{aligned}$$

$$\begin{aligned}
 & \|\tilde{u}\|_{H,(0 \rightarrow k+1),\delta}^2 \\
 &= \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^{k+1} \|a_{m\ell}\|_{H,0,\delta}^2 \\
 &\leq C \left( \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^{k-1} (\|b_{m\ell}\|_{H,0,\delta-2}^2 + \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} [1 + \ell(\ell + 1)]^{k+1/2} |c_{m\ell}|^2) \right) \\
 &= C \left( \|F\|_{\mathcal{A}_{H\delta-2}^{(0,k-1)}}^2 + \|h\|_{H^{k+1/2}(S^2)}^2 \right)
 \end{aligned}$$

It then follows that



$$\|\tilde{u}\|_{\mathcal{A}_H^{(2,k+1)}} = \max \left\{ \|u\|_{H,(2 \rightarrow k-1),\delta}, \|u\|_{H,(1 \rightarrow k),\delta}, \|u\|_{H,(0 \rightarrow k+1),\delta} \right\} \quad (3.14)$$

$$\leq C \left( \|F\|_{\mathcal{A}_H^{(0,k-1)}}^2 + \|h\|_{H^{k+1/2}(S^2)}^2 \right) \quad (3.15)$$

as needed.

The rest of the proof is then devoted to prove estimates **H-Est** and **C-Est**.

We first introduce a piece of notation. Given 2 quantities  $\alpha, \beta$ , we will write  $\alpha \lesssim \beta$  if there exists a constant  $C > 0$  depending only on  $n, m_0$  and  $\delta$ , such that  $\alpha \leq C\beta$ . In particular, the constant will not depend on  $\tilde{u}, F, h, m$ , and  $\ell$ . In this notation, the estimates that we will be proving are

$$\begin{aligned} \|a''_{m\ell}\|_{H,0,\delta-2}^2 + [1 + \ell(\ell + 1)] \|a'_{m\ell}(r)\|_{H,0,\delta-1}^2 + [1 + \ell(\ell + 1)]^2 \|a_{m\ell}(r)\|_{H,0,\delta}^2 \\ \lesssim \|b_{m\ell}\|_{H,0,\delta-2}^2 + [1 + \ell(\ell + 1)]^{3/2} |c_{m\ell}|^2 \end{aligned} \quad (\mathbf{H-Est})$$

$$\begin{aligned} \|a''_{m\ell}\|_{C,0,\delta-2}^2 + [1 + \ell(\ell + 1)] \|a'_{m\ell}(r)\|_{C,0,\delta-1}^2 + [1 + \ell(\ell + 1)]^2 \|a_{m\ell}(r)\|_{C,0,\delta}^2 \\ \lesssim \|b_{m\ell}\|_{C,0,\delta-2}^2 + [1 + \ell(\ell + 1)]^2 |c_{m\ell}|^2 \end{aligned} \quad (\mathbf{C-Est})$$

for every  $m$  and  $\ell$ .

The relation between  $\tilde{u}, F$  and  $h$ , namely  $F = \Delta_{g_{sc}} \tilde{u}$  and  $h = \tilde{u}(nm_0)$ , imply the following differential equation on the coefficients  $a_{m\ell}, b_{m\ell}$ , and  $c_{m\ell}$ :

$$\begin{cases} r(r - 2m_0)a''_{m\ell}(r) + 2(r - m_0)a'_{m\ell}(r) - \ell(\ell + 1)a_{m\ell}(r) = r(r - 2m_0)b_{m\ell}(r), & r \in [nm_0, \infty) \\ a_{m\ell}(nm_0) = c_{m\ell} \end{cases} \quad (3.16)$$

We first consider the case  $\ell = 0$ . we integrate once the above differential equation to get

$$r(r - 2m_0)a'_{00}(r) = \int_{nm_0}^r s(s - 2m_0)b_{00}(s)ds + n(n - 2)m_0^2 a'_{00}(nm_0) \quad (3.17)$$

which immediately gives the following estimate on  $\|a'_{00}\|_{C,0,\delta-1}$ :

$$\|a'_{00}\|_{C,0,\delta-1} \lesssim \|b_{00}\|_{C,0,\delta-2} + |a'_{00}(nm_0)| \quad (3.18)$$

To estimate  $\|a'_{00}\|_{H,0,\delta-1}$ , we use equation (3.17) to get

$$\int_{nm_0}^{\infty} r^{-2\delta+1} (a'_{00}(r))^2 dr \lesssim \int_{nm_0}^{\infty} \frac{r^{-2\delta+1}}{r^2(r-2m_0)^2} \left( \int_{nm_0}^r s(s-2m_0)b_{00}(s)ds \right)^2 dr + |a'_{00}(nm_0)|^2 \quad (3.19)$$

$$\lesssim \int_{nm_0}^{\infty} r^{-2\delta+3} (b_{00}(r))^2 dr + |a'_{00}(nm_0)|^2 \quad (3.20)$$

where Hardy's inequality was used in the last line. We then conclude the following estimate on  $\|a'_{00}\|_{H,0,\delta-1}$ :

$$\|a'_{00}\|_{H,0,\delta-1} \lesssim \|b_{00}\|_{H,0,\delta-2} + |a'_{00}(nm_0)| \quad (3.21)$$

We divide equation (3.17) by  $r(r-2m_0)$  and integrate to get

$$a_{00}(r) = c_{00} + \int_{nm_0}^r \frac{1}{s(s-2m_0)} \int_{nm_0}^s s'(s'-2m_0)b_{00}(s')ds'ds + a'_{00}(nm_0) \ln \left( \frac{n(r-2m_0)}{(n-2)r} \right) \frac{n(n-2)m_0}{2} \quad (3.22)$$

which, in the same manner as for  $a'_{00}$ , gives the following estimates:

$$\|a_{00}\|_{C,0,\delta} \lesssim |c_{00}| + \|b_{00}\|_{C,0,\delta-2} + |a'_{00}(nm_0)| \quad (3.23)$$

$$\|a_{00}\|_{H,0,\delta} \lesssim |c_{00}| + \|b_{00}\|_{H,0,\delta-2} + |a'_{00}(nm_0)| \quad (3.24)$$

Using the fact that  $a_{00}$  vanishes at infinity, we take the limit as  $r$  goes to infinity in equation (3.22) to get the following expression for  $a'_{00}(nm_0)$  in terms of  $c_{00}$  and  $b_{00}$ :

$$a'_{00}(nm_0) = \frac{2}{n(n-2) \ln \left( \frac{n}{n-2} \right)} \left( -c_{00} - \int_{nm_0}^{\infty} \frac{1}{r(r-2m_0)} \int_{nm_0}^r s(s-2m_0)b_{00}(s)dsdr \right) \quad (3.25)$$

which allows us to estimate  $a'_{00}(nm_0)$  to get

$$|a'_{00}(nm_0)| \lesssim |c_{00}| + \|b_{00}\|_{H,0,\delta-2} \quad (3.26)$$

$$|a'_{00}(nm_0)| \lesssim |c_{00}| + \|b_{00}\|_{C,0,\delta-2} \quad (3.27)$$

We now get an estimate for  $a''_{00}(r)$ . Using the ODE in (3.16) we isolate for  $a''_{00}(r)$  to get:

$$a''_{00}(r) = b_{00}(r) - \frac{2(r-m_0)}{r(r-2m_0)}a'_{00}(r) \quad (3.28)$$

It then follows that

$$\|a''_{00}\|_{C,0,\delta-2} \lesssim \|b_{00}\|_{C,0,\delta-2} + \|a'_{00}\|_{C,0,\delta-1} \quad (3.29)$$

$$\|a''_{00}\|_{H,0,\delta-2} \lesssim \|b_{00}\|_{H,0,\delta-2} + \|a'_{00}\|_{H,0,\delta-1} \quad (3.30)$$

Combining the estimates for  $a_{00}$ ,  $a'_{00}$  and  $a''_{00}$  in equations (3.23), (3.24), (3.18), (3.21), (3.29) and (3.30) together with the estimates for  $|a'_{00}(nm_0)|$  in equations (3.26) and (3.27), we finally deduce the desired estimates:

$$\|a_{00}\|_{H,2,\delta} \lesssim |c_{00}| + \|b_{00}\|_{H,0,\delta-2} \quad (3.31)$$

$$\|a_{00}\|_{C,2,\delta} \lesssim |c_{00}| + \|b_{00}\|_{C,0,\delta-2} \quad (3.32)$$

We now deal with the case  $\ell \geq 1$ .

*Proving H-Est for  $\ell \geq 1$*

We multiply both sides of the differential equation in (3.16) by  $r^{-2\delta-1}a_{m\ell}(r)$  and integrate by parts to obtain:

$$\begin{aligned} & \int_{nm_0}^{\infty} r^{-2\delta-1}r(r-2m_0)a_{m\ell}^2(r)dr + \int_{nm_0}^{\infty} \left[ \ell(\ell+1) + (2\delta+1) \left( \delta - \frac{m_0(2\delta+1)}{r} \right) \right] r^{-2\delta-1}a_{m\ell}^2(r)dr \\ &= - \int_{nm_0}^{\infty} r^{-2\delta-1}a_{m\ell}(r)b_{m\ell}(r)dr - (nm_0)^{-2\delta}m_0(n-2)c_{m\ell}a'_{m\ell}(nm_0) + \frac{-2\delta-1}{2}(nm_0)^{-2\delta-1}m_0(n-2)c_{m\ell}^2 \end{aligned} \quad (3.33)$$

We observe that for  $\ell \geq 1$ ,

$$\ell(\ell+1) + (2\delta+1) \left( \delta - \frac{m_0(2\delta+1)}{r} \right) > \ell(\ell+1) - \frac{1}{2} \quad (3.34)$$

and, hence, we have that

$$\begin{aligned} \|a'_{m\ell}\|_{H,0,\delta-1}^2 + [1 + \ell(\ell + 1)] \|a_{m\ell}\|_{H,0,\delta}^2 &\lesssim \int_{nm_0}^{\infty} r^{-2\delta-1} r(r - 2m_0) a_{m\ell}^{\prime 2}(r) dr \\ &\quad + \int_{nm_0}^{\infty} \left[ \ell(\ell + 1) + (2\delta + 1) \left( \delta - \frac{m_0(2\delta + 1)}{r} \right) \right] r^{-2\delta-1} a_{m\ell}^2(r) dr \end{aligned} \quad (3.35)$$

Deriving an upper bound for the expression in right hand side of equation (3.33) will then lead to an upper bound for  $\|a'_{m\ell}\|_{H,0,\delta-1}^2 + [1 + \ell(\ell + 1)] \|a_{m\ell}\|_{H,0,\delta}^2$ .

We obtain an estimate for  $a'_{m\ell}(nm_0)$  by multiplying equation (3.16) by  $a'_{m\ell}(r)$  and integrating by parts to get

$$\int_{nm_0}^{\infty} (r - m_0) a_{m\ell}^{\prime 2}(r) dr + \ell(\ell + 1) \frac{1}{2} c_{m\ell}^2 - \frac{1}{2} nm_0^2 (n - 2) a_{m\ell}^{\prime 2}(nm_0) = \int_{nm_0}^{\infty} r(r - 2m_0) a'_{m\ell}(r) b_{m\ell}(r) dr \quad (3.36)$$

where we used the fact that  $a_{m\ell}^{\prime 2}(r)r(r - 2m_0) = o(1)$  and  $a_{m\ell}^2(r) = o(1)$ . By estimating the right side of the above equation as follows

$$\left| \int_{nm_0}^{\infty} r(r - 2m_0) a'_{m\ell}(r) b_{m\ell}(r) dr \right| \lesssim [\ell(\ell + 1)]^{1/2} \|a'_{m\ell}\|_{H,0,\delta-1}^2 + [\ell(\ell + 1)]^{-1/2} \|b_{m\ell}\|_{H,0,\delta-2}^2, \quad (3.37)$$

we deduce that

$$|a'_{m\ell}(nm_0)| \lesssim [\ell(\ell + 1)]^{1/4} \|a'_{m\ell}\|_{H,0,\delta-1} + [\ell(\ell + 1)]^{-1/4} \|b_{m\ell}\|_{H,0,\delta-2} + [\ell(\ell + 1)]^{1/2} |c_{m\ell}| \quad (3.38)$$

We can now estimate the right hand side of equation (3.33) to get

$$\begin{aligned} & - \int_{nm_0}^{\infty} r^{-2\delta-1} a_{m\ell}(r) b_{m\ell}(r) dr - (nm_0)^{-2\delta} m_0 (n - 2) c_{m\ell} a'_{m\ell}(nm_0) + \frac{-2\delta - 1}{2} (nm_0)^{-2\delta-1} m_0 (n - 2) c_{m\ell}^2 \\ & \lesssim \|a_{m\ell}\|_{H,0,\delta} \|b_{m\ell}\|_{H,0,\delta-2} + |c_{m\ell}| \left( [\ell(\ell + 1)]^{1/4} \|a'_{m\ell}\|_{H,0,\delta-1} + [\ell(\ell + 1)]^{-1/4} \|b_{m\ell}\|_{H,0,\delta-2} + [\ell(\ell + 1)]^{1/2} |c_{m\ell}| \right) \\ & \quad + c_{m\ell}^2 \end{aligned} \quad (3.39)$$

We estimate each term appearing in the right hand side of the above equation. Let  $\varepsilon > 0$  that will be chosen to be small later on. Then we have

$$\|a_{m\ell}\|_{H,0,\delta} \|b_{m\ell}\|_{H,0,\delta-2} \leq \epsilon \ell(\ell+1) \|a_{m\ell}\|_{H,0,\delta}^2 + \frac{D(\epsilon)}{\ell(\ell+1)} \|b_{m\ell}\|_{H,0,\delta-2}^2 \quad (3.40)$$

$$c_{m\ell} \|b_{m\ell}\|_{H,0,\delta-2} \leq \frac{[\ell(\ell+1)]^{3/4}}{2} c_{m\ell}^2 + \frac{1}{2[\ell(\ell+1)]^{3/4}} \|b_{m\ell}\|_{H,0,\delta-2}^2 \quad (3.41)$$

$$c_{m\ell} \|a'_{m\ell}\|_{H,0,\delta-1} \leq \epsilon [\ell(\ell+1)]^{-1/4} \|a'_{m\ell}\|_{H,0,\delta-1}^2 + D(\epsilon) [\ell(\ell+1)]^{1/4} c_{m\ell}^2 \quad (3.42)$$

where  $D = D(\epsilon)$  is a constant depending on  $\epsilon$ .

Combining the above, we get

$$\begin{aligned} & \|a'_{m\ell}\|_{H,0,\delta-1}^2 + [1 + \ell(\ell+1)] \|a_{m\ell}\|_{H,0,\delta}^2 \quad (3.43) \\ & \leq C\epsilon \left( \ell(\ell+1) \|a_{m\ell}\|_{H,0,\delta}^2 + \|a'_{m\ell}\|_{H,0,\delta-1}^2 \right) + CD(\epsilon) \left( [\ell(\ell+1)]^{1/2} c_{m\ell}^2 + \frac{1}{\ell(\ell+1)} \|b_{m\ell}\|_{H,0,\delta-2}^2 \right) \quad (3.44) \end{aligned}$$

for some constant  $C > 0$  that only depends on  $n$ ,  $m_0$ , and  $\delta$ . Choosing  $\epsilon$  to be small, we can absorb the expression multiplied to  $C\epsilon$  in the above equation to the left hand side to finally deduce

$$\|a'_{m\ell}\|_{H,0,\delta-1}^2 + [1 + \ell(\ell+1)] \|a_{m\ell}\|_{H,0,\delta}^2 \lesssim [1 + \ell(\ell+1)]^{-1} \|b_{m\ell}\|_{H,0,\delta-2}^2 + [1 + \ell(\ell+1)]^{1/2} c_{m\ell}^2 \quad (3.45)$$

which is the desired estimate for  $\|a'_{m\ell}\|_{H,0,\delta-1}$  and  $\|a_{m\ell}\|_{H,0,\delta}$ .

What is left is to estimate  $\|a''_{m\ell}\|_{H,0,\delta-2}$ . We use the ODE in (3.16) to get

$$r^{-2\delta-1} r^2 (r-2m_0)^2 (a''_{m\ell}(r))^2 = r^{-2\delta-1} \left[ b_{m\ell}(r) + \ell(\ell+1)a_{m\ell}(r) - 2(r-m_0)a'_{m\ell}(r) \right]^2 \quad (3.46)$$

We can then estimate

$$\begin{aligned} \|a''_{m\ell}\|_{H,0,\delta-2}^2 &\lesssim \|b_{m\ell}\|_{H,0,\delta-2}^2 + [1 + \ell(\ell + 1)]^2 \|a_{m\ell}\|_{H,0,\delta}^2 + \|a'_{m\ell}\|_{H,0,\delta-1}^2 + \ell(\ell + 1) \|b_{m\ell}\|_{H,0,\delta-2} \|a_{m\ell}\|_{H,0,\delta} \\ &\quad + \|a'_{m\ell}\|_{H,0,\delta-1} \|b_{m\ell}\|_{H,0,\delta-2} + \ell(\ell + 1) \|a_{m\ell}\|_{H,0,\delta} \|a'_{m\ell}\|_{H,0,\delta-1} \end{aligned} \quad (3.47)$$

$$\lesssim [1 + \ell(\ell + 1)] \|a'_{m\ell}\|_{H,0,\delta-1}^2 + [1 + \ell(\ell + 1)]^2 \|a_{m\ell}\|_{H,0,\delta}^2 + \|b_{m\ell}\|_{H,0,\delta-2}^2 \quad (3.48)$$

$$\lesssim [1 + \ell(\ell + 1)]^{3/2} c_{m\ell}^2 + \|b_{m\ell}\|_{H,0,\delta-2}^2 \quad (3.49)$$

where we used equation (3.45) in the last line. Combining the above equation with equation (3.45), we finally get the desired estimate **H-Est**:

$$\|a''_{m\ell}\|_{H,0,\delta-2}^2 + [1 + \ell(\ell + 1)] \|a'_{m\ell}\|_{H,0,\delta-1}^2 + [1 + \ell(\ell + 1)]^2 \|a_{m\ell}\|_{H,0,\delta}^2 \lesssim \|b_{m\ell}\|_{H,0,\delta-2}^2 + [1 + \ell(\ell + 1)]^{3/2} c_{m\ell}^2 \quad (3.50)$$

*Proving C-Est for  $\ell \geq 1$*

For  $r \in [nm_0, \infty)$  define

$$z := \frac{r}{m_0} - 1, \quad R := n - 1, \quad h_{m\ell}(z) := a_{m\ell}(r), \quad f_{m\ell}(z) := r(r - 2m_0)b_{m\ell}(r)$$

Note that  $R > 1$  since  $n > 2$ . The desired estimate in **C-Est** in terms of  $h$  is then

$$\|h''_{m\ell}\|_{C,0,\delta-2}^2 + [1 + \ell(\ell + 1)] \|h'_{m\ell}\|_{C,0,\delta-1}^2 + [1 + \ell(\ell + 1)]^2 \|h_{m\ell}\|_{C,0,\delta}^2 \lesssim \|f_{m\ell}\|_{C,0,\delta}^2 + [1 + \ell(\ell + 1)]^2 c_{m\ell}^2 \quad (\mathbf{C-Est'})$$

The IVP for  $a_{m\ell}$  in (3.16) becomes

$$\begin{cases} (z^2 - 1)h''_{m\ell}(z) + 2zh'_{m\ell}(z) - \ell(\ell + 1)h_{m\ell}(z) = f_{m\ell}(z), & z \in [R, \infty) \\ h_{m\ell}(R) = c_{m\ell} \\ h_{m\ell} \in H_\delta^2([R, \infty)) \cap C_\delta^2([R, \infty)) \end{cases} \quad (3.51)$$

The above ODE is the Legendre differential equation; the Legendre functions of the first and second kind,  $P_\ell$  and  $Q_\ell$ , are two linearly independent solutions to the homogeneous equation in (3.51) (i.e. with  $f_{m\ell} = 0$ ) satisfying the following asymptotics as  $z \rightarrow \infty$  (see [OR14] chapter 5 section 12):

$$P_\ell(z) = O(z^\ell), \quad Q_\ell(z) = O(z^{-\ell-1}) \quad (3.52)$$

We will frequently use some properties of those functions discussed and proved in section A.3 in the Appendix.

We normalize  $P_\ell$  and  $Q_\ell$  so that

$$\lim_{z \rightarrow \infty} z^{-\ell} P_\ell(z) = 1, \quad \lim_{z \rightarrow \infty} z^{\ell+1} Q_\ell(z) = 1 \quad (3.53)$$

Using the method of Frobenius, we can expand  $P_\ell$  and  $Q_\ell$  as a sum of powers of  $z$  on  $[R, \infty)$ , which we present in proposition A.4 in the Appendix. We rewrite the proposition here for convenience.

**Proposition 3.4.**  *$P_\ell$  and  $Q_\ell$  admit an expansion of the following form. For  $z > 1$ ,*

$$P_\ell(z) = \sum_{k=0}^{\ell} a_k z^{\ell-k}, \quad Q_\ell(z) = \sum_{k=0}^{\infty} b_k z^{-\ell-1-k} \quad (3.54)$$

where the coefficients  $a_k$  and  $b_k$  are defined recursively as follows:

$$a_0 = b_0 = 1, \quad a_1 = b_1 = 0$$

for  $k \geq 2$ , 
$$a_k = \frac{(\ell - k + 2)(\ell - k + 1)}{k^2 - k(2\ell + 1)} a_{k-2}, \quad b_k = \frac{(\ell + k - 1)(\ell + k)}{k(2\ell + k + 1)} b_{k-2}$$

We observe immediately from the above that  $Q_\ell(z)$  is positive and  $z^{\ell+1} Q_\ell(z)$  is decreasing on  $[R, \infty)$ .

Using the variation of parameters method (see [BO13]), we can explicitly write the solution to (3.51):

$$h_{m\ell}(z) = A Q_\ell(z) + P_\ell(z) \int_z^\infty Q_\ell(t) f_{m\ell}(t) [W(t)(t^2 - 1)]^{-1} dt + Q_\ell(z) \int_R^z P_\ell(t) f_{m\ell}(t) [W(t)(t^2 - 1)]^{-1} dt \quad (3.55)$$

where  $W(t) := P_\ell(t) Q'_\ell(t) - P'_\ell(t) Q_\ell(t)$  is the Wronskian and  $A$  is defined by

$$A = \frac{1}{Q_\ell(R)} \left( c_{m\ell} - P_\ell(R) \int_R^\infty Q_\ell(t) f_{m\ell}(t) [W(t)(t^2 - 1)]^{-1} dt \right) \quad (3.56)$$

Note that  $W(t)(t^2 - 1)$  is a constant by Lagrange's identity (see [Rico2] pg 354). We compute that constant to be  $2\ell + 1$  by taking the limit as  $z$  goes to  $\infty$  in the expansion of  $P_\ell$  and  $Q_\ell$ .

We summarize here some estimates on the Legendre functions uniform in  $z$  and  $\ell$  that we prove in the Appendix (check proposition A.5): There exists a constant  $C = C(R)$  such that for any  $\ell \geq 1$  and  $z \in [R, \infty)$ , the following holds

$$z^{-\ell}|P_\ell(z)| \leq C \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^{-\ell}, \quad z^{\ell+1}|Q_\ell(z)| \leq C \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^\ell \quad (3.57)$$

$$z^{-(\ell-1)}|P'_\ell(z)| \leq C\ell \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^{-\ell}, \quad z^{\ell+2}|Q'_\ell(z)| \leq C\ell \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^\ell \quad (3.58)$$

We note that the function  $\frac{2z}{z + \sqrt{z^2 - 1}}$  is decreasing on  $[R, \infty)$  and is bounded below and above by 1 and 2 respectively. Using the expression for  $h_{m\ell}$  in equation (3.55) and the uniform bounds of  $P_\ell$  and  $Q_\ell$  in equation (3.57), we obtain

$$z^{-\delta}|h_{m\ell}(z)| \leq z^{-\delta}|A||Q_\ell(z)| + z^{-\delta}\frac{1}{2\ell+1}|P_\ell(z)| \int_z^\infty |Q_\ell(t)||f_{m\ell}(t)|dt + z^{-\delta}\frac{1}{2\ell+1}|Q_\ell(z)| \int_R^z |P_\ell(t)||f_{m\ell}(t)|dt \quad (3.59)$$

$$\begin{aligned} &\leq z^{-\delta}|A|Q_\ell(R)R^{\ell+1}z^{-\ell-1} \\ &\quad + \frac{C^2}{2\ell+1} \left( \sup_{t \geq R} t^{-\delta}|f_{m\ell}(t)| \right) z^{\ell-\delta} \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^{-\ell} \int_z^\infty \left( \frac{2t}{t + \sqrt{t^2 - 1}} \right)^\ell t^{-\ell-1}t^\delta dt \\ &\quad + \frac{C^2}{2\ell+1} \left( \sup_{t \geq R} t^{-\delta}|f_{m\ell}(t)| \right) z^{-\ell-1-\delta} \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^\ell \int_R^z \left( \frac{2t}{t + \sqrt{t^2 - 1}} \right)^{-\ell} t^\ell t^\delta dt \end{aligned} \quad (3.60)$$

$$\leq |A|Q_\ell(R)R^{-\delta} + \frac{C^2}{2\ell+1} \left( \sup_{t \geq R} t^{-\delta}|f_{m\ell}(t)| \right) \left( \frac{1}{\ell-\delta} + \frac{1}{\ell+1+\delta} \left( 1 - \left( \frac{R}{z} \right)^{\ell+\delta+1} \right) \right) \quad (3.61)$$

It immediately follows that

$$\|h_{m\ell}\|_{C,0,\delta} \lesssim |A|Q_\ell(R) + \ell^{-2} \|f_{m\ell}\|_{C,0,\delta} \quad (3.62)$$

To derive an estimate for  $|A|Q_\ell(R)$ , we use equation (3.56) to get



$$|A|Q_\ell(R) \leq |c_{m\ell}| + \frac{C^2}{2\ell+1} \left( \sup_{t \geq R} t^{-\delta} |f_{m\ell}(t)| \right) R^\ell \left( \frac{2R}{R + \sqrt{R^2 - 1}} \right)^{-\ell} \int_R^\infty \left( \frac{2t}{t + \sqrt{t^2 - 1}} \right)^\ell t^{-\ell-1} t^\delta dt \quad (3.63)$$

$$\leq |c_{m\ell}| + \frac{C^2 R^\delta}{(2\ell+1)(\ell-\delta)} \left( \sup_{t \geq R} t^{-\delta} |f_{m\ell}(t)| \right) \quad (3.64)$$

The above estimate for  $|A|Q_\ell(R)$  together with equation (3.62) implies

$$\|h_{m\ell}\|_{C_{0,\delta}} \lesssim |c_{m\ell}| + \ell^{-2} \|f_{m\ell}\|_{C_{0,\delta}} \quad (3.65)$$

To achieve an estimate for  $\|h'_{m\ell}\|_{C_{0,\delta-1}}$ , we take the derivative of equation (3.55) to get

$$h'_{m\ell}(z) = A Q'_\ell(z) + \frac{1}{2\ell+1} P'_\ell(z) \int_z^\infty Q_\ell(t) f_{m\ell}(t) dt + \frac{1}{2\ell+1} Q'_\ell(z) \int_R^z P_\ell(t) f_{m\ell}(t) dt \quad (3.66)$$

In a similar manner, we will apply the uniform bounds on  $P_\ell$  and  $Q_\ell$  in equations (3.57) and (3.58) to obtain the desired estimate for  $\|h'_{m\ell}\|_{C_{0,\delta-1}}$ . Using the above equation for  $h'_{m\ell}$  as well as equations (3.57) and (3.58), we get

$$\begin{aligned} z^{-\delta+1} |h'_{m\ell}(z)| &\leq z^{-\delta+1} |A| |Q'_\ell(z)| \\ &\quad + z^{-\delta+1} \frac{1}{2\ell+1} |P'_\ell(z)| \int_z^\infty |Q_\ell(t)| |f_{m\ell}(t)| dt + z^{-\delta+1} \frac{1}{2\ell+1} |Q'_\ell(z)| \int_R^z |P_\ell(t)| |f_{m\ell}(t)| dt \\ &\leq z^{-\delta+1} |A| |Q'_\ell(z)| \\ &\quad + \frac{C^2 \ell}{2\ell+1} \left( \sup_{t \geq R} t^{-\delta} |f_{m\ell}(t)| \right) z^{\ell-\delta} \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^{-\ell} \int_z^\infty \left( \frac{2t}{t + \sqrt{t^2 - 1}} \right)^\ell t^{-\ell-1} t^\delta dt \\ &\quad + \frac{C^2 \ell}{2\ell+1} \left( \sup_{t \geq R} t^{-\delta} |f_{m\ell}(t)| \right) z^{-\ell-1-\delta} \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^\ell \int_R^z \left( \frac{2t}{t + \sqrt{t^2 - 1}} \right)^{-\ell} t^\ell t^\delta dt \\ &\leq z^{-\delta+1} |A| |Q'_\ell(z)| + \frac{C^2 \ell}{2\ell+1} \left( \sup_{t \geq R} t^{-\delta} |f_{m\ell}(t)| \right) \left( \frac{1}{\ell-\delta} + \frac{1}{\ell+1+\delta} \left( 1 - \left( \frac{R}{z} \right)^{\ell+\delta+1} \right) \right) \end{aligned} \quad (3.67)$$

$$(3.68)$$

$$(3.69)$$

We estimate the term  $z^{-\delta+1}|A||Q'_\ell(z)|$ . First, we observe from the expansion of  $Q_\ell(z)$  in proposition A.4 that  $z^{\ell+2}Q'_\ell(z)$  is negative and increasing on  $[R, \infty)$ , which in particular implies that

$$z^{-\delta+1}|A||Q'_\ell(z)| \leq z^{-\delta+1}|A||Q'_\ell(R)|R^{\ell+2}z^{-\ell-2} \quad (3.70)$$

$$\leq z^{-\delta-1} \left(\frac{R}{z}\right)^\ell R^2|A||Q'_\ell(R)| \quad (3.71)$$

$$\leq R^{-\delta+1}|A||Q'_\ell(R)| \quad (3.72)$$

We then use the recursive relation for  $Q_\ell$  in equation (A.44) to deduce that

$$|Q'_\ell(R)| = \frac{\ell}{R^2-1}(-RQ_\ell(R) + Q_{\ell-1}(R)) \quad (3.73)$$

$$< \frac{\ell}{R^2-1}Q_{\ell-1}(R) \quad (3.74)$$

which, in light of equations (3.72) and (3.64), implies that

$$z^{-\delta-1}|A||Q'_\ell(z)| \leq \frac{\ell R^{-\delta+1}}{R^2-1}|A|Q_{\ell-1}(R) \quad (3.75)$$

$$\leq \frac{\ell R^{-\delta+1}}{R^2-1} \left( |c_{m\ell}| + \frac{C^2 R^\delta}{(2\ell-1)(\ell-1-\delta)} \left( \sup_{t \geq R} t^{-\delta} |f_{m\ell}(t)| \right) \right) \quad (3.76)$$

The above together with equation (3.69) finally lead to the desired estimate for  $\|h'_{m\ell}\|_{C,0,\delta-1}$ :

$$\|h'_{m\ell}\|_{C,0,\delta-1} \lesssim \ell |c_{m\ell}| + \ell^{-1} \|f_{m\ell}\|_{C,0,\delta} \quad (3.77)$$

What is left is estimating  $\|h''_{m\ell}\|_{C,0,\delta-2}$ . Using the ODE satisfied by  $h_{m\ell}$  in (3.51), we have

$$z^{2-\delta}h''_{m\ell}(z) = -2\frac{z^{3-\delta}}{z^2-1}h'_{m\ell}(z) + \frac{\ell(\ell+1)z^{2-\delta}}{z^2-1}h_{m\ell}(z) + \frac{z^{2-\delta}}{z^2-1}f_{m\ell}(z) \quad (3.78)$$

which then, using equations (3.65) and (3.77) implies

$$\|h''_{m\ell}\|_{C,0,\delta-2} \lesssim \|h'_{m\ell}\|_{C,0,\delta-1} + \ell(\ell+1) \|h_{m\ell}\|_{C,0,\delta} + \|f_{m\ell}\|_{C,0,\delta} \quad (3.79)$$

$$\lesssim \ell^2 |c_{m\ell}| + \|f_{m\ell}\|_{C,0,\delta} \quad (3.80)$$

The above equation together with equations (3.65) and (3.77) finally imply the desired estimate:

$$\|h''_{m\ell}\|_{C,0,\delta-2}^2 + [1 + \ell(\ell + 1)] \|h'_{m\ell}\|_{C,0,\delta-1}^2 + [1 + \ell(\ell + 1)]^2 \|h_{m\ell}\|_{C,0,\delta}^2 \lesssim \|f_{m\ell}\|_{C,0,\delta}^2 + [1 + \ell(\ell + 1)]^2 c_{m\ell}^2$$

(C-Est')

This concludes the proof of the lemma. □

## REDUCTION OF THE PROBLEM

---

In this section, we reduce the static Einstein vacuum equations into a simpler system involving ODEs, the Laplace equation on  $M$ , and first order partial differential equations on  $\partial M$ .

Let  $g$  be a metric on  $M$  of the form  $dr^2 + g(r)$ , where  $g(r)$  is a metric on  $S^2$  for each  $r \in [nm_0, \infty)$ . The level sets of the function  $r$  defines a foliation with leaves denoted by  $S_r$ . We define the unit vector field  $n := \frac{\partial}{\partial r}$  that is normal to the foliation. We denote by  $\nabla$  and  $\text{div}$  the covariant derivative and divergence with respect to the induced metric  $g(r)$  on  $S_r$ , and  $d$  the exterior derivative on  $S_r$ .

We then define the second fundamental form  $K$  as the  $(0,2)$  symmetric tensor field on  $M$  that is tangential to the leaves  $S_r$  of the foliation and satisfies:

$$K(X, Y) := g(\nabla_n X, Y) \quad (4.1)$$

for vector fields  $X, Y$  on  $M$  that are tangential to  $S_r$ . We will decompose  $K$  into the sum of its traceless and trace parts:

$$K = \hat{K} + \frac{1}{2} \text{tr}K g$$

Note that  $\text{Hess}_g(r) = K$  and  $\Delta_g r = \text{tr}K$ .

The following equations on the leaves  $S_r$  describe the evolution of the geometry on  $M$  in terms of  $K$  and  $\text{Ric}$  (see [CK93] and [KN12]). Given coordinates  $(r, \theta^1, \theta^2)$  on  $M$ ,

$$\partial_r \text{tr}K + \frac{1}{2} (\text{tr}K)^2 + |\hat{K}|^2 = -R_{00} \quad (4.2)$$

$$\nabla_r \hat{K}_{ij} + \text{tr}K \hat{K}_{ij} = - \left[ R_{ij} + \frac{1}{2} g_{ij} (R_{00} - R) \right] \quad (4.3)$$

$$R_{S_r} - \frac{1}{2}(\text{tr}K)^2 + |\hat{K}|^2 = R - 2R_{00} \quad (4.4)$$

$$\nabla^j \hat{K}_{ji} - \frac{1}{2} \nabla_i \text{tr}K = R_{0i} \quad (4.5)$$

$$\partial_r g_{ij} = 2\hat{K}^{ij} + \text{tr}K g_{ij} \quad (4.6)$$

where  $i, j = 1, 2$ ,  $R_{S_r}$  is the scalar curvature of  $(S_r, g(r))$ ,  $R$  is the scalar curvature on  $(M, g)$ , and  $\nabla$  is the connection on  $(S_r, g(r))$ . Moreover,  $R_{00} := \text{Ric}(n, n)$  and  $R_{0i} := \text{Ric}(n, \frac{\partial}{\partial \theta^i})$  for  $i = 1, 2$ .

We note that the left side of equation (4.3) can be simplified as follows

$$\nabla_r \hat{K}_{ij} + \text{tr}K \hat{K}_{ij} = \partial_r \hat{K}_{ij} - 2\Gamma_{0j}^l \hat{K}_{il} + \text{tr}K \hat{K}_{ij} \quad (4.7)$$

$$= \partial_r \hat{K}_{ij} \quad (4.8)$$

Equation (4.3) can then be written as follows:

$$(\mathcal{L}_{\frac{\partial}{\partial r}} \hat{K})_{ij} = - \left[ R_{ij} + \frac{1}{2} g_{ij} (R_{00} - R) \right] \quad (4.9)$$

The above equations determine all the components of the Ricci curvature of  $g$ . More specifically, if the right hand sides of equations (4.2) to (4.5) are known on all the leaves, then the Ricci curvature can be fully recovered. In fact, if we in addition know that  $(g, u)$  solves the conformal static vacuum Einstein equations for some function  $u$  on  $M$ , then, due to the contracted Bianchi identities, equations (4.4) and (4.5) need only to be imposed on the boundary for the Ricci curvature to be fully recovered. The next proposition will prove this fact and will demonstrate the desired reduction of our problem.

**Reduction Theorem.** *Let  $(\gamma_{\mathfrak{B}}, \frac{1}{2} \text{tr}K_{\mathfrak{B}})$  be Bartnik data. Let  $g = dr^2 + g(r)$  and  $u$  be a metric and function on  $M$  respectively, where  $g(r)$  is a metric on  $S^2$  for every  $r \in [nm_0, \infty)$ . The pair  $(g, f) = (e^{-2u} g, e^u)$  solves the static Einstein vacuum equations with Bartnik data  $(\gamma_{\mathfrak{B}}, \frac{1}{2} \text{tr}K_{\mathfrak{B}})$  if and only if  $(g, u)$  satisfies*

$$\Delta_g u = 0, \quad \text{on } M \quad (4.10)$$

$$\partial_r \text{tr} K + \frac{1}{2} \text{tr} K^2 + |\hat{K}|^2 + 2(\partial_r u)^2 = 0, \quad \text{on } M \quad (4.11)$$

$$\mathcal{L}_{\frac{\partial}{\partial r}} \hat{K} + [2\mathcal{d}u \otimes \mathcal{d}u + g(r) ((\partial_r u)^2 - |\nabla u|^2)] = 0, \quad \text{on } M \quad (4.12)$$

$$2|\nabla u|^2 - 2(\partial_r u)^2 - |\hat{K}|^2 - R_{\partial M} + \frac{1}{2} \text{tr} K^2 = 0, \quad \text{on } \partial M \quad (4.13)$$

$$2(\partial_r u)\mathcal{d}u - \text{div}(\hat{K}) + \frac{1}{2} \mathcal{d} \text{tr} K = 0, \quad \text{on } \partial M \quad (4.14)$$

$$e^{-2u} g|_{\partial M} = \gamma_{\mathfrak{B}}, \quad \text{on } \partial M \quad (4.15)$$

$$e^u (\text{tr} K|_{\partial M} - 2\partial_r u) = \text{tr} K_{\mathfrak{B}}, \quad \text{on } \partial M \quad (4.16)$$

*Proof.* The “only if ” direction is clear from equations (4.2) to (4.5). We prove the “if ” direction.

Suppose  $(g, u)$  satisfy equations (4.10) to (4.16). It suffices to show that  $\text{Ric} = 2\mathcal{d}u \otimes \mathcal{d}u$ . We first decompose the Ricci curvature of  $g$  with respect to the foliation. Let  $\Pi$  be the  $(1, 1)$  projection tensor field defined by

$$\Pi_{\nu}^{\mu} = \delta_{\nu}^{\mu} - n^{\mu} n_{\nu} \quad (4.17)$$

We then define the function  $Q$ , the 1-form  $P$  tangential to the foliation, and the  $(0, 2)$  symmetric tensor field  $S$  tangential to the foliation as follows:

$$Q := \text{Ric}(n, n), \quad P_{\mu} := \Pi_{\mu}^{\mu'} n^{\nu} R_{\mu'\nu}, \quad S_{\mu\nu} := \Pi_{\mu}^{\mu'} \Pi_{\nu}^{\nu'} R_{\mu'\nu'} \quad (4.18)$$

The Ricci curvature of  $g$  can then be written in the following way:

$$\text{Ric} = Q\underline{n} \otimes \underline{n} + P \otimes \underline{n} + \underline{n} \otimes P + S \quad (4.19)$$

where  $\underline{n}$  is the 1-form achieved by lowering the index for  $n$ . We will omit the underbar when we write  $\underline{n}$  in components.

Define the function  $H$  on  $M$  and the 1-form  $A$  tangent to the foliation in the following way:

$$H := R - 2|\nabla u|^2 \quad (4.20)$$

$$A := P - 2n(u)\mathcal{d}u \quad (4.21)$$

We now compare equations (4.2) - (4.5) with equations (4.11) - (4.14). From Equation (4.2) and (4.11), we deduce on  $M$  that

$$Q = 2n(u)^2 \quad (4.22)$$

From equation (4.9) and (4.12), we deduce on  $M$  that

$$S = 2\mathcal{d}u \otimes \mathcal{d}u + \frac{1}{2}\gamma H \quad (4.23)$$

We also have by definition of  $A$ :

$$P = 2n(u)\mathcal{d}u + A \quad (4.24)$$

From equation (4.4) and (4.13), it follows that on  $\partial M$ ,

$$R - 2Q = 2|\nabla u|^2 - 4n(u)^2 \quad (4.25)$$

which gives us:

$$H|_{\partial M} = 0 \quad (4.26)$$

From equation (4.5) and (4.14), we get:

$$A|_{\partial M} = 0$$

To prove the statement, we just need to show that  $H, A = 0$ .

We first prove the following lemma.

**Lemma 4.1.** *Let  $dr^2 + g(r)$  be a metric on  $M$  where  $g(r)$  is a metric on  $S^2$  for every  $r \in [nm_0, \infty)$ . Suppose that the Ricci decomposition relative to the foliation defined by  $r$ , as written in equation (4.19), is*

$$Q = 2n(u)^2, \quad P = 2n(u)\mathcal{d}u + A, \quad S = 2\mathcal{d}u \otimes \mathcal{d}u + \frac{1}{2}g(r)H \quad (4.27)$$

where  $u$  is a harmonic function on  $M$ ,  $H$  is a function, and  $A$  is a 1-form tangent to the foliation.

Then  $A$  and  $H$  satisfy

$$(\nabla_n A)_k + A_i K_k^i + \text{tr} K A_k = 0 \quad (4.28)$$

$$\nabla_n H + H \text{tr} K = 2 \text{div}(A) \quad (4.29)$$

*Proof.* Recall the second Bianchi identity:

$$\frac{1}{2}\nabla_\nu R = \nabla^\mu R_{\nu\mu} \quad (4.30)$$

where  $\mu, \nu = 0, 1, 2$ .

We can write the Ricci curvature as follows:

$$Ric = Q\underline{n} \otimes \underline{n} + P \otimes \underline{n} + \underline{n} \otimes P + S \quad (4.31)$$

We compute the divergence of the tensor  $2du \otimes du$  to be:

$$\nabla^\mu (2du \otimes du)_{\mu\nu} = 2\nabla^\mu du_\mu du_\nu + 2du_\mu \nabla^\mu du_\nu \quad (4.32)$$

$$= \Delta u du_\nu + 2\text{Hess}(u)(\nabla u, \partial_\nu) \quad (4.33)$$

$$= \nabla_\nu |\nabla u|^2 \quad (4.34)$$

where  $\Delta u = 0$  was used in the last line. Using the Bianchi identities, we get

$$\frac{1}{2}\nabla_\nu R = \nabla^\mu \left( 2du \otimes du + A \otimes \underline{n} + \underline{n} \otimes A + \frac{1}{2}H\gamma \right)_{\mu\nu} \quad (4.35)$$

$$= \nabla^\mu (2du \otimes du)_{\mu\nu} + \nabla^\mu A_\mu n_\nu + A_\mu \nabla^\mu n_\nu + \nabla^\mu n_\mu A_\nu + n_\mu \nabla^\mu A_\nu \quad (4.36)$$

$$+ \frac{1}{2}\nabla_{\nu'} H \Pi \Pi_{\nu'}^{\nu'} + \frac{1}{2}H \nabla^\mu (g_{\mu\nu} - n_\mu n_\nu) \quad (4.37)$$

$$= \nabla_\nu |\nabla u|^2 + \nabla^\mu A_\mu n_\nu + A_\mu K_\nu^\mu + \text{tr}K A_\nu + \nabla_n A_\nu$$

$$+ \frac{1}{2}\nabla_{\nu'} H \Pi \Pi_{\nu'}^{\nu'} - \frac{1}{2}H \text{tr}K n_\nu$$

where  $\Pi_\mu^{\nu'} = \delta_\mu^{\nu'} - n^{\nu'} n_\mu$ . We also used the fact that  $\gamma_{\mu\nu} = g_{\mu'\nu'} \Pi_\mu^{\mu'} \Pi_\nu^{\nu'} = g_{\mu\nu} - n_\mu n_\nu$ .

Using fermi coordinates  $(r, \theta^1, \theta^2)$  and letting  $\nu = i = 1, 2$ , we get

$$(\nabla_n A)_i + A_j K_i^j + \text{tr}K A_i = 0 \quad (4.38)$$

Letting  $\nu = 0$ , we get

$$\nabla_n H + H \text{tr}K = 2\nabla^\mu A_\mu \quad (4.39)$$

as desired.



□

We then have that  $A$  satisfies,

$$\begin{cases} (\nabla_n A)_k + A_i K_k^i + \text{tr} K A_k = 0, & \text{on } M \\ A = 0, & \text{on } \partial M \end{cases} \quad (4.40)$$

By the existence and uniqueness theory of ODEs, it follows that  $A = 0$ .

Since  $\text{div}(A) = 0$ , we get that  $H$  satisfies

$$\begin{cases} \nabla_n H + H \text{tr} K = 0, & \text{on } M \\ H = 0, & \text{on } \partial M \end{cases} \quad (4.41)$$

By invoking again the existence and uniqueness theorem of ODEs, we deduce that  $H = 0$ . □

## PROOF OF THE MAIN THEOREM

---

The reduction theorem in section 4 suggests that we study the map

$$\begin{aligned}
\Psi : & \mathcal{M}^{k+1}(\partial M) \times H^k(\partial M) \times \mathcal{M}_\delta^k(M) \times \mathcal{A}_\delta^{(2,k+1)} \\
\rightarrow & \mathcal{A}_{\delta-2}^{(0,k-1)}(M) \times L_{\delta-2}^2\left([nm_0, \infty); H^k(S^2)\right) \times L_{\delta-2}^2\left([nm_0, \infty); \mathcal{H}^k(S^2)\right) \\
& \times H^{k-1}(\partial M) \times \Omega^{k-1}(\partial M) \times \mathcal{H}^k(\partial M) \times H^k(\partial M)
\end{aligned}$$

$$\Psi(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}}, g, u) := \begin{pmatrix} \Delta_g u \\ \partial_r trK + \frac{1}{2}trK^2 + |\hat{K}|^2 + 2(\partial_r u)^2 \\ \mathcal{L}_{\frac{\partial}{\partial r}} \hat{K} + [2\mathfrak{d}u \otimes \mathfrak{d}u + g(r)((\partial_r u)^2 - |\nabla u|^2)] \\ 2|\nabla u|^2 - 2(\partial_r u)^2 - |\hat{K}|^2 - R_{\partial M} + \frac{1}{2}trK^2 \\ 2(\partial_r u)\mathfrak{d}u - \mathfrak{d}\hat{\sigma}(\hat{K}) + \frac{1}{2}\mathfrak{d}trK \\ e^{-2u}g|_{\partial M} - \gamma_{\mathfrak{B}} \\ trK|_{\partial M} - e^{-u}(trK_{\mathfrak{B}} + 2e^u\partial_r u) \end{pmatrix} \tag{5.1}$$

where  $trK$  and  $\hat{K}$  are with respect to the metric  $g$ , and  $R_{\partial M}$  is with respect to the metric  $e^{2u}\gamma_{\mathfrak{B}}$ . Furthermore, norms  $|\cdot|$  used in the second, third and fourth line are with respect to the metric  $g$ .

We wish to show that there exists a map taking Bartnik data  $(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}})$  close to Schwarzschild data to a pair  $(g, u)$  satisfying  $\Psi(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}}, g, u) = 0$ , showing that  $(g, f) = (e^{-2u}g, e^u)$  solves the static Einstein vacuum equations with Bartnik data  $(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}})$ . This can be achieved by first attempting to show that the linearization of  $\Psi$  with respect to  $(g, u)$  at  $(\gamma_{g_{sc}}, \frac{1}{2}trK_{g_{sc}}, g_{sc}, u_{sc})$  is an isomorphism, or merely surjective, and then

invoking the implicit function theorem. However, the linearization of the contracted Codazzi equation, in the fifth line of the definition of  $\Psi$ , leads to obstructions to surjectivity stemming from the divergence operator acting on symmetric traceless tensors on  $S^2$ . More specifically, we are faced with the cokernel of the divergence operator: a 6-dimensional space of obstructions equal to the space of conformal Killing vector fields on  $S^2$ .

This difficulty does not preclude the possibility of finding solutions given arbitrary Bartnik data close to Schwarzschild data. A similar situation arises when one attempts to show the existence of metrics on the sphere with prescribed scalar curvature (see [KW75]). The operator of study will not satisfy the conditions for the inverse function theorem, yet existence holds as shown in [AH21]. In our case, we circumvent this difficulty by introducing an artificial object, in the form of a vector field  $X$ , to the meaning of a solution to our problem, proving its existence using the implicit function theorem, and then finally showing that this vector field  $X$  vanishes, yielding a solution to the original problem.

### 5.1 DEFINITION OF THE ARTIFICIAL VECTOR FIELD $X$

As explained in the introduction of section 5, the contracted Codazzi equations give rise to obstructions that are in correspondence with the space of conformal Killing vector fields on  $S^2$ . We will overcome these seeming obstructions by introducing an artificial vector field  $X$  to the definition of a solution; this means that the solution will consist of a metric  $g$ , a function  $u$ , and a vector field  $X$ . This needs to be done in a way so that, firstly, the corresponding modified problem is solvable, and secondly, the artificial vector field, in fact, vanishes for a solution to the modified problem, yielding a solution to the original problem. To achieve this, the artificial vector field  $X$  needs to be carefully defined, which will require a certain way of uniquely extending conformal Killing fields from  $S^2$  to the ambient manifold  $M$ . This procedure will be outlined in this section.

Notably, Huang and An have also introduced an artificial vector field  $X$  in [AH21] and [AH22b] for analogous purposes; specifically, they define  $X$  to be a vector field that vanishes on the boundary and asymptotically approaches a Killing vector field at infinity. In contrast, we will define  $X$  to be a vector field that satisfies  $\left(\mathcal{L}_{\frac{\partial}{\partial r}} X\right)^T = 0$  on the boundary and asymptotically approaches a conformal killing vector field on  $(M, g_{sc})$  that restricts to a conformal Killing vector field on  $(\partial M, \gamma_{S^2})$ .

Given a metric  $g$  and a vector field  $X$  on  $M$ , we denote by  $\mathcal{L}_{g,conf}X$  and  $\Delta_{g,conf}X$  the conformal Lie derivative of  $g$  with respect to  $X$  and the conformal laplacian of  $X$  defined by

$$\mathcal{L}_{g,conf}X := \widehat{\mathcal{L}_X g}, \quad \Delta_{g,conf}X := \operatorname{div}_g(\mathcal{L}_{g,conf}X) \quad (5.2)$$

where  $\widehat{\mathcal{L}_X g}$  is the traceless part of  $\mathcal{L}_X g$ . It follows that  $X$  is conformal Killing on  $(M, g)$  if and only if  $\mathcal{L}_{g,conf}X = 0$ .

**Definition 5.1.** Given a conformal Killing vector field  $X_{CK}$  on  $(\partial M, \gamma_{S^2})$ , we denote by  $\overline{X_{CK}}$  the unique vector field on  $(M, g_{sc})$  extending  $X_{CK}$  on  $\partial M$  and satisfying the evolution equation

$$\mathcal{L}_{\frac{\partial}{\partial r}} \overline{X_{CK}} = 0, \quad \text{in } M \quad (5.3)$$

Also, we will use “ $\operatorname{div}_{\gamma_{S^2}}(X_{CK})$ ” to denote both the divergence of  $X_{CK}$  on  $(\partial M, \gamma_{S^2})$  and the same function extended to a function on  $M$  independent of  $r$ . It should be clear from context which one we are referring to.

**Definition 5.2.** Define the space  $\mathcal{X}_\infty$  as the space of conformal killing vector fields  $X_\infty$  on  $(M, g_{sc})$  of the form

$$X_\infty = f(r) \left( \operatorname{div}_{\gamma_{S^2}}(X_{CK}) \right) \frac{\partial}{\partial r} + h(r) \overline{X_{CK}} \quad (5.4)$$

where  $f = f(r)$  and  $h = h(r)$  are smooth functions on  $M$  such that  $f = 0$  and  $h = 1$  on  $\partial M$  and  $X_{CK}$  is a conformal Killing vector field on  $(\partial M, \gamma_{S^2})$ .

In the case that  $X_{CK}$  is Killing, equation (5.4) becomes

$$X_\infty = h(r) \overline{X_{CK}} \quad (5.5)$$

**Lemma 5.3.** Let  $X_{CK}$  be a nontrivial conformal Killing vector field on  $(\partial M, \gamma_{S^2})$ .

- (a) Suppose  $X_{CK}$  is Killing on  $(\partial M, \gamma_{S^2})$ . Then  $h \equiv 1$  is the unique smooth function  $h = h(r)$  on  $M$  in which  $h = 1$  on  $\partial M$  and  $X_\infty$ , defined by equation (5.5), is conformal Killing on  $(M, g_{sc})$ . In fact,  $X_\infty$  would also be Killing.
- (b) Suppose  $X_{CK}$  is not Killing on  $(\partial M, \gamma_{S^2})$ . There exists unique smooth functions  $f = f(r)$  and  $h = h(r)$  on  $M$  such that  $f \equiv 0$ ,  $h \equiv 1$  on  $\partial M$  and the vector field  $X_\infty$  defined by (5.4) is conformal Killing on  $(M, g_{sc})$ . Furthermore,  $f = O(r^2)$  and  $h = O(r^2)$ .

In particular,  $\mathcal{X}_\infty$  is a 6 dimensional vector space of conformal Killing vector fields on  $(M, g_{sc})$ .

*Proof.* We will repeatedly use the following identity of the Lie derivative: for any vector fields  $X, Y, Z$  and any  $(0, 2)$  tensor field  $T$  on  $M$ ,

$$[\mathcal{L}_X T](Y, Z) = X\left(T(Y, Z)\right) - T([X, Y], Z) - T(Y, [X, Z]) \quad (5.6)$$

We first prove (a). Suppose  $X_{CK}$  is Killing on  $(\partial M, \gamma_{S^2})$ . It is clear that  $h(r)\overline{X_{CK}}$  is Killing on  $(M, g_{sc})$  for  $h \equiv 1$  on  $M$  as it is a rotation vector on the spherically symmetric Schwarzschild manifold. Now suppose  $h = h(r)$  is a smooth function such that  $h \equiv 1$  on  $\partial M$  and  $h(r)\overline{X_{CK}}$  is conformal Killing on  $M$ . In particular, we have that

$$0 = [\mathcal{L}_{h(r)\overline{X_{CK}}} g_{sc}] \left( \frac{\partial}{\partial r}, \overline{X_{CK}} \right) \quad (5.7)$$

$$= g_{sc} \left( \left[ \frac{\partial}{\partial r}, h(r)\overline{X_{CK}} \right], \overline{X_{CK}} \right) + g_{sc} \left( [Y, h(r)\overline{X_{CK}}], \frac{\partial}{\partial r} \right) \quad (5.8)$$

$$= h'(r)g_{sc}(\overline{X_{CK}}, \overline{X_{CK}}) \quad (5.9)$$

In view of the fact that  $X_{CK} \neq 0$ , it follows that  $h' \equiv 0$  and so  $h \equiv 1$  on  $M$  as needed.

We now prove (b). Suppose  $X_{CK}$  is not Killing on  $(\partial M, \gamma_{S^2})$ . Let  $f = f(r)$  and  $h = h(r)$  be smooth functions on  $M$ . Recall that the metric  $g_{sc}$  can be written as

$$g_{sc} = dr^2 + r(r - 2m_0)\gamma_{S^2}$$

Only for the proof of this lemma, we will denote the function  $\mathcal{L}_{X_{CK}} \gamma_{S^2}$  by  $B_{X_{CK}}$  for simplicity of the notation. Recall that  $B_{X_{CK}}$  is understood as a function on  $M$  or a function on  $\partial M$  depending on the context, and that  $B_{X_{CK}}$  as a function on  $M$  is constant in  $r$ .

Since  $X_{CK}$  is not Killing, we have that  $B_{X_{CK}}$  is nonzero. Moreover, after fixing a spherical coordinate system on  $\partial M$ , the vector field  $X_{CK}$  can be written as a linear combination of the vector fields  $\nabla_{\gamma_{S^2}} Y_m^{\ell=1}$ , for  $m = -1, 0, 1$ , where  $Y_m^{\ell=1}$  are the  $\ell = 1$  spherical harmonics on  $\partial M$ . In particular, it holds that

$$\nabla_{\gamma_{S^2}} B_{X_{CK}} = -2X_{CK} \quad (5.10)$$

This implies that

$$\nabla_{g_{sc}} (B_{X_{CK}}) = \frac{1}{r(r-2m_0)} \nabla_{\gamma_{S^2}} (B_{X_{CK}}) \quad (5.11)$$

$$= -\frac{2}{r(r-2m_0)} \overline{X_{CK}} \quad (5.12)$$

For  $X_\infty := f(r)B_{X_{CK}}\frac{\partial}{\partial r} + h(r)\overline{X_{CK}}$  and arbitrary vector fields  $Y, Z$  tangent to the foliation, we compute

$$[\mathcal{L}_{X_\infty} g_{sc}] \left( \frac{\partial}{\partial r}, \frac{\partial}{\partial r} \right) = 2g_{sc} \left( \left[ \frac{\partial}{\partial r}, X_\infty \right], \frac{\partial}{\partial r} \right) \quad (5.13)$$

$$= 2f'(r)B_{X_{CK}} \quad (5.14)$$

$$[\mathcal{L}_{X_\infty} g_{sc}] \left( \frac{\partial}{\partial r}, Y \right) = g_{sc} \left( \left[ \frac{\partial}{\partial r}, X_\infty \right], Y \right) + g_{sc} \left( [Y, X_\infty], \frac{\partial}{\partial r} \right) \quad (5.15)$$

$$= h'(r)g_{sc}(\overline{X_{CK}}, Y) + f(r)Y(B_{X_{CK}}) \quad (5.16)$$

$$= h'(r)g_{sc}(\overline{X_{CK}}, Y) + f(r)g_{sc}(\nabla B_{X_{CK}}, Y) \quad (5.17)$$

$$= \left( h'(r) - \frac{2}{r(r-2m_0)}f(r) \right) g_{sc}(\overline{X_{CK}}, Y) \quad (5.18)$$

$$[\mathcal{L}_{X_\infty} g_{sc}] (Y, Z) = [\mathcal{L}_{f(r)B_{X_{CK}}\frac{\partial}{\partial r}} g_{sc}] (Y, Z) + [\mathcal{L}_{h(r)\overline{X_{CK}}} g_{sc}] (Y, Z) \quad (5.19)$$

$$= f(r) B_{X_{CK}} \operatorname{tr} K_{sc} g_{sc}(Y, Z) + r(r-2m_0)h(r) [\mathcal{L}_{X_{CK}} \gamma_{S^2}] (Y, Z) \quad (5.20)$$

$$= f(r) B_{X_{CK}} \operatorname{tr} K_{sc} g_{sc}(Y, Z) + r(r-2m_0)h(r) B_{X_{CK}} \gamma_{S^2}(Y, Z) \quad (5.21)$$

$$= [f(r)\operatorname{tr} K_{sc} + h(r)] B_{X_{CK}} g_{sc}(Y, Z) \quad (5.22)$$

In the above calculation, we used the fact that  $X_{CK}$  is conformal Killing on each leaf and hence  $\mathcal{L}_{X_{CK}} \gamma_{S^2} = B_{X_{CK}} \gamma_{S^2}$ .

It follows that  $\mathcal{L}_{X_\infty} g_{sc}$  is conformal to  $g_{sc}$ ,  $f \equiv 0$  and  $h \equiv 1$  on  $\partial M$  if and only if the pair  $(f, h)$  satisfy the following on  $[nm_0, \infty)$ .

$$\begin{cases} 2f'(r) = f(r)\text{tr}K_{sc} + h(r), \\ h'(r) = \frac{2}{r(r-2m_0)}f(r), \\ f(nm_0) = 0, \\ h(nm_0) = 1 \end{cases} \quad (5.23)$$

This decouples to the following initial value problems for  $f$  and  $h$  on  $[nm_0, \infty)$ .

$$\begin{cases} f''(r) - \frac{(r-m_0)}{r(r-2m_0)}f'(r) + \frac{2m_0^2}{r^2(r-2m_0)^2}f(r) = 0 \\ f(nm_0) = 0 \\ f'(nm_0) = \frac{1}{2} \end{cases} \quad (5.24)$$

$$\begin{cases} h'(r) = \frac{2}{r(r-2m_0)}f(r) \\ h(nm_0) = 1 \end{cases} \quad (5.25)$$

Invoking the existence and uniqueness theorem for ODEs, it follows that there exists unique smooth functions  $f$  and  $g$  satisfying the above initial value problems.

We have then proven that there exists unique smooth functions  $f = f(r)$  and  $h = h(r)$  on  $M$  such that  $f \equiv 0$ ,  $h \equiv 1$  on  $\partial M$  and the vector field  $X_\infty$  is conformal Killing on  $(M, g_{sc})$ .

We utilize Fuchsian theory to establish that both  $f$  and  $h$  are  $O(r^2)$ . We first observe that the ODE for  $f$  in equation (5.24) has a regular singular point at infinity. We can then express  $f$  as a Frobenius series as follows (see [BO13]):

$$f(r) = (r - m_0)^\alpha \sum_{n=0}^{\infty} a_n (r - m_0)^{-n} \quad (5.26)$$

where  $\alpha \in \mathbb{R}$  is to be determined and  $a_0 \neq 0$ . We substitute this expression of  $f$  into the ODE in (5.24) to get

$$\begin{aligned} & \sum_{n=0}^{\infty} a_n (\alpha - n)(\alpha - n - 1)(r - m_0)^{\alpha - n - 2} - \sum_{n=0}^{\infty} a_n (\alpha - n)(r - m_0)^{\alpha - n - 2} \left(1 + \frac{m_0^2}{r(r - 2m_0)}\right) \\ & + 2m_0^2 \sum_{n=0}^{\infty} a_n (r - m_0)^{\alpha - n - 4} \frac{(r - m_0)^4}{r^2(r - 2m_0)^2} \end{aligned} = 0 \quad (5.27)$$

Upon examining the highest power of  $r - m_0$ , we deduce that  $\alpha$  must satisfy the equation

$$\alpha(\alpha - 1) - \alpha = 0 \tag{5.28}$$

implying that  $\alpha$  can only be 0 or 2. It follows that  $f = O(r^2)$ . The fact that  $h = O(r^2)$  follows immediately from equation (5.25). □

*Remark 5.4.* Note that any vector field  $X_\infty$  in  $\mathcal{X}_\infty$  satisfies

$$\left( \mathcal{L}_{\frac{\partial}{\partial r}} X_\infty \right)^T = 0, \quad \text{on } \partial M$$

Hence, the lemma proves an existence and uniqueness result for an overdetermined problem: for any conformal Killing vector field  $X_{CK}$  on  $(\partial M, \gamma_{S^2})$ , there exists a unique conformal Killing vector field  $X_\infty$  on  $(M, g_{sc})$  satisfying the following boundary conditions on  $\partial M$ :

$$X_\infty|_{\partial M} = X_{CK}, \quad \left( \mathcal{L}_{\frac{\partial}{\partial r}} X_\infty \right)^T = 0 \tag{5.29}$$

Furthermore,  $X_\infty$  will be of the form as in equation (5.4) for some functions  $f = f(r)$  and  $h = h(r)$ .

**Definition 5.5.** We define  $\widehat{\mathcal{X}}_\delta^2(M)$  to be all vector fields  $X \in \mathcal{X}_\delta^2(M)$  (see definition 2.5) such that  $X|_{\partial M}$  is tangent to  $\partial M$  and  $\left( \mathcal{L}_{\frac{\partial}{\partial r}} X \right)^T = 0$  on  $\partial M$ . The artificial vector field  $X$  will be chosen to live in the space  $\widehat{\mathcal{X}}_\delta^2(M) \oplus \mathcal{X}_\infty(M)$ . The reasons for this choice will be clear in the next sections.

## 5.2 DEFINITION AND EXISTENCE OF THE MODIFIED SOLUTION

In this section, we will define the modified problem and its solutions, which we call "the modified solutions", and prove their existence. Here and onwards, we fix a number  $\delta \in (-1, -\frac{1}{2}]$  and an integer  $k \geq 5$ .

Let  $\eta$  be a smooth cut off function on  $[nm_0, \infty)$  satisfying  $\eta(r) = 1$  for  $r \geq nm_0 + 2$  and  $\eta(r) = 0$  for  $r \leq nm_0 + 1$ . Given  $X \in \widehat{\mathcal{X}}_\delta^2(M) \oplus \mathcal{X}_\infty(M)$ , define the function  $F(X)$  on  $M$  by

$$F(X) := e^{-r^4 \eta(r) |X|^2} \tag{5.30}$$

where  $|\cdot|$  is taken with respect to  $g_{sc}$ .



Given  $g = dr^2 + g(r) \in \mathcal{M}_\delta^k(M)$  and  $X_\infty \in \mathcal{X}_\infty$ , define  $\omega(g, X_\infty)$  to be the 1-form on  $\partial M$  achieved by lowering the index of  $X_\infty|_{\partial M}$  with respect to  $g(nm_0)$ . Note that  $X_\infty|_{\partial M}$  is a conformal Killing field on  $(\partial M, \gamma_{S^2})$  by definition of the space  $\mathcal{X}_\infty$ .

Define  $\Phi$  by:

$$\begin{aligned} \Phi : & \mathcal{M}^{k+1}(\partial M) \times H^k(\partial M) \times \mathcal{M}_\delta^k(M) \times \mathcal{A}_\delta^{(2,k+1)}(M) \times \left( \widehat{\mathcal{X}}_\delta^2(M) \oplus \mathcal{X}_\infty(M) \right) \\ \rightarrow & \mathcal{A}_{\delta-2}^{(0,k-1)}(M) \times L_{\delta-2}^2\left([nm_0, \infty); H^k(S^2)\right) \times L_{\delta-2}^2\left([nm_0, \infty); \mathcal{H}^k(S^2)\right) \times \mathcal{X}_{\delta-2}^0(M) \\ & \times H^{k-1}(\partial M) \times \Omega^{k-1}(\partial M) \times \mathcal{H}^k(\partial M) \times H^k(\partial M) \end{aligned}$$

$$\Phi(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}}, g, u, X) := \left( \begin{array}{ll} \Delta_g u & \text{on } M \\ \partial_r trK + \frac{1}{2}trK^2 + |\hat{K}|^2 + 2(\partial_r u)^2 & \text{on } M \\ \mathcal{L}_{\frac{\partial}{\partial r}} \hat{K} + [2\mathcal{A}u \otimes \mathcal{A}u + g(r)((\partial_r u)^2 - |\nabla u|^2)] & \text{on } M \\ \Delta_{g,conf}(F(X)X) & \text{on } M \\ 2|\nabla u|^2 - 2(\partial_r u)^2 - |\hat{K}|^2 - R_{\partial M} + \frac{1}{2}trK^2 & \text{on } \partial M \\ 2(\partial_r u)\mathcal{A}u - \mathcal{A}\hat{K} + \frac{1}{2}\mathcal{A}trK + \omega(g, X_\infty) & \text{on } \partial M \\ e^{-2u}g|_{\partial M} - \gamma_{\mathfrak{B}} & \text{on } \partial M \\ trK|_{\partial M} - e^{-u}(trK_{\mathfrak{B}} + 2e^u\partial_r u) & \text{on } \partial M \end{array} \right) \quad (5.31)$$

where  $X_\infty$  is the projection of  $X$  onto  $\mathcal{X}_\infty(M)$ ,  $trK$  and  $\hat{K}$  are with respect to the metric  $g$ , and  $R_{\partial M}$  is with respect to the metric  $e^{2u}\gamma_{\mathfrak{B}}$ . Furthermore, norms  $|\cdot|$  used in the second, third and fifth equation are with respect to the metric  $g$ .

**Definition 5.6.** Given Bartnik data  $(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}})$ , we say that a 3-tuple  $(g, u, X)$  is a modified solution if  $\Phi(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}}, g, u, X) = 0$ .

*Remark 5.7.* In view of proposition , a modified solution is a solution to the original problem if and only if  $X = 0$ .

The main tool to obtain the existence of the modified problem is the implicit function theorem on Banach manifolds (see [AMR12]), which is stated here for convenience.

**Theorem 5.8.** *Let  $U \subset E$ ,  $V \subset F$  be open subsets of Banach spaces  $E$  and  $F$ , and let  $\Psi : U \times V \rightarrow G$  be a  $C^r$  map to a Banach space  $G$ , with  $r \geq 1$ . For some  $x_0 \in U$ ,  $y_0 \in V$ , assume the partial derivatives in the second argument  $D_2\Psi(x_0, y_0) : F \rightarrow G$  is an isomorphism. Then there are neighbourhoods  $U_0$  of  $x_0$  and  $W_0$  of  $\Psi(x_0, y_0)$  and a unique  $C^r$  map  $H : U_0 \times W_0 \rightarrow V$  such that for all  $(x, w) \in U_0 \times W_0$ ,  $\Psi(x, H(x, w)) = w$ .*

The map  $\Phi$  is indeed  $C^1$  near  $(\gamma_{g_{sc}}, \frac{1}{2}trK_{g_{sc}}, g_{sc}, u_{sc}, 0)$ . To see this, we first note the following:

- The map  $u \mapsto du \otimes du$  is  $C^1$  from  $\mathcal{A}_\delta^{(2,k+1)}(M)$  to  $H_{2\delta-2}^1([nm_0, \infty); \mathcal{H}^k(S^2))$ .
- The map  $g \mapsto trK$  is  $C^1$  from  $\mathcal{M}_\delta^k(M)$  to  $H_{\delta-1}^1([nm_0, \infty); H^k(S^2))$ .
- The map  $g \mapsto \hat{K}$  is  $C^1$  from  $\mathcal{M}_\delta^k(M)$  to  $H_{\delta-1}^1([nm_0, \infty); \mathcal{H}^k(S^2))$ .

This immediately shows that each line, excluding the fourth line, in the definition of  $\Phi$  is  $C^1$ . It remains to show that the map  $(g, X) \mapsto \Delta_{g,conf}(F(X)X)$  is  $C^1$  from  $\mathcal{M}_\delta^k(M) \times \left( \hat{\mathcal{X}}_\delta^2(M) \oplus \mathcal{X}_\infty(M) \right)$  to  $\mathcal{X}_{\delta-2}^0$  near  $(g_{sc}, 0)$ . This follows directly from the smoothness of  $F(X)$  and the following identity of the conformal laplacian (see [Yor74]):

$$\Delta_{g,conf}(F(X)X)_\mu = \Delta_g(F(X)X)_\mu + \frac{1}{3}\nabla_\mu \left( \operatorname{div}_g(F(X)X) \right) + R_{\mu\nu}F(X)X^\nu \quad (5.32)$$

We can then differentiate  $\Phi$  at  $(\gamma_{g_{sc}}, \frac{1}{2}trK_{g_{sc}}, g_{sc}, u_{sc}, 0)$  and study its derivative.

Let  $D\Phi_{sc}$  denote the derivative of  $\Phi$  with respect to the last three components evaluated at  $(\gamma_{g_{sc}}, \frac{1}{2}trK_{g_{sc}}, g_{sc}, u_{sc}, 0)$  where

$$\begin{aligned} D\Phi_{sc} : & T_{g_{sc}}\mathcal{M}_\delta^k \times \mathcal{A}_\delta^{(2,k+1)} \times \left( \hat{\mathcal{X}}_\delta^2(M) \oplus \mathcal{X}_\infty(M) \right) \\ \rightarrow & \mathcal{A}_{\delta-2}^{(0,k-1)}(M) \times L_{\delta-2}^2 \left( [nm_0, \infty); H^k(S^2) \right) \times L_{\delta-2}^2 \left( [nm_0, \infty); \mathcal{H}^k(S^2) \right) \\ & \times H^{k-1}(\partial M) \times \Omega^{k-1}(\partial M) \times \mathcal{H}^k(\partial M) \times H^k(\partial M) \end{aligned}$$

**Proposition 5.9.**  $D\Phi_{sc}$  is an isomorphism.

*Proof.* The proof of this will be the content of section (5.4).  $\square$

We can now conclude the existence theorem for the extended problem.

**Theorem 5.10.** *There exists a neighbourhood  $\mathcal{U}$  of  $(\gamma_{g_{sc}}, \frac{1}{2}trK_{g_{sc}})$  in  $\mathcal{M}^{k+1}(\partial M) \times H^k(\partial M)$  and a unique  $\mathcal{C}^1$  map  $\mathbf{H} : (\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}}) \mapsto (g, u, X)$  on  $\mathcal{U}$  into  $\mathcal{M}_{\delta}^k(M) \times \mathcal{A}_{\delta}^{(2,k+1)}(M) \times \widehat{\mathcal{X}}_{\delta}^2(M) \oplus \mathcal{X}_{\infty}(M)$  satisfying*

$$\Phi(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}}, \mathbf{H}(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}})) = 0, \quad \text{for all } (\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}}) \in \mathcal{U} \quad (5.33)$$

*Proof.* Follows from proposition (5.9) and the implicit function theorem on Banach manifolds.  $\square$

### 5.3 THE VANISHING OF $X$ FOR MODIFIED SOLUTIONS $(g, u, X)$

The next step is to show that if  $(g, u, X)$  is a modified solution, then  $X = 0$ , yielding a solution  $(g, u)$  to the conformal static vacuum Einstein equations.

Let  $(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}}) \in \mathcal{U}$  be Bartnik data and let  $(g, u, X) := \mathbf{H}(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}})$  be the corresponding modified solution.

We first find the Ricci curvature of the metric  $g$ .

**Proposition 5.11.** *The Ricci decomposition of  $g = dr^2 + g(r)$  relative to the foliation defined by  $r$  is given by*

$$Ric = Q\underline{n} \otimes \underline{n} + P \otimes \underline{n} + \underline{n} \otimes P + S \quad (5.34)$$

$$Q = 2n(u)^2, \quad P = 2n(u)\underline{d}u + A, \quad S = 2\underline{d}u \otimes \underline{d}u + \frac{1}{2}g(r)H \quad (5.35)$$

where  $H$  and  $A$  are the unique function on  $M$  and 1-form on  $M$  tangent to the foliation satisfying:

$$\begin{cases} \nabla_n A_k + A_i K_k^i + trK A_k = 0, & \text{on } M \\ A = \omega(g, X_{\infty}), & \text{on } \partial M \end{cases} \quad (5.36)$$

$$\begin{cases} \nabla_n H + H \operatorname{tr} K = 2 \operatorname{div}(A), & \text{on } M \\ H = 0, & \text{on } \partial M \end{cases} \quad (5.37)$$

*Proof.* Lemma (4.1) shows that  $A$  and  $H$  satisfy the desired transport equations on  $M$ . The boundary condition for  $A$  and  $H$  follow by comparing equations

$$2|\nabla u|^2 - 2(\partial_r u)^2 - |\hat{K}|^2 - R_{\partial M} + \frac{1}{2} \operatorname{tr} K^2 = 0, \quad \text{and} \quad 2(\partial_r u) \not{d}u - \not{d}\hat{v}(\hat{K}) + \frac{1}{2} \not{d}\operatorname{tr} K + \omega(g, X_\infty) = 0$$

with equations (4.4) and (4.5).  $\square$

The relation between the Ricci curvature of  $g$  and  $u$  as described in the above proposition leads to the following regularity result.

**Proposition 5.12.** *The following holds for any modified solution  $(g, u, X)$ .*

- *The Ricci curvature of  $g$  is  $\mathcal{C}^1$  away from the boundary. Furthermore, there exists a universal constant  $C > 0$  such that for  $R > nm_0$ ,*

$$\sup_{r>R} r^2 (|\operatorname{Ric}| + r|\nabla \operatorname{Ric}|) \leq C \|\gamma_\infty - \gamma_{S^2}\|_{\mathcal{H}^k(\partial M)} + o(R^\delta) \quad (5.38)$$

as  $R$  goes to  $\infty$ .

- *The vector field  $F(X)X$  lies in  $\mathcal{X}_\delta^3(M)$ .*

*Proof.* We first find explicit expressions for  $A$  and  $H$ . Letting  $(r, \theta^1, \theta^2)$  be fermi coordinates, we compute for  $i = 1, 2$ ,

$$\nabla_n A_i = \partial_r A_i - \Gamma_{0i}^j A_j \quad (5.39)$$

$$= \partial_r A_i - K_i^j A_j \quad (5.40)$$

Equation (5.36) for  $A$  then becomes

$$\begin{cases} \partial_r A_i + \operatorname{tr} K A_i = 0, & \text{on } M \\ A = \omega(g, X_\infty), & \text{on } \partial M \end{cases} \quad (5.41)$$

which gives

$$A_i(r) = \frac{1}{L(r)} \omega_i \quad (5.42)$$

where  $L(r) := \exp\left(\int_{nm_0}^r \text{tr}K(s)ds\right)$ .

We then solve equation(5.37) for  $H$  to obtain

$$H(r) = \frac{1}{L(r)} \int_{nm_0}^r L(s)2\text{div}(A(s))ds \quad (5.43)$$

From proposition (2.10), we have

$$\left| \text{tr}K - \frac{2}{r} \right| \in H_{\delta-1}^1([nm_0, \infty); H^k(S^2)), \quad \hat{K} \in H_{\delta-1}^1([nm_0, \infty); \mathcal{H}^k(S^2)) \quad (5.44)$$

It then follows that  $A \in H_{loc}^2([nm_0, \infty); \mathcal{H}^k(S^2))$  and  $\text{div}(A) \in H_{loc}^1([nm_0, \infty); \mathcal{H}^{k-1}(S^2))$ , which in turn implies that  $H \in H_{loc}^2([nm_0, \infty); H^{k-1}(S^2))$ .

Furthermore, since  $k \geq 5$ , the Sobolev embedding described in proposition (2.10) imply that  $\text{tr}K$  and  $\hat{K}$  are continuous in  $r$ , have 3 continuous angular derivatives, and satisfy

$$\text{tr}K = \frac{2}{r} + o(r^{\delta-1}), \quad |\hat{K}| = o(r^{\delta-1}) \quad (5.45)$$

Using the asymptotics of  $\text{tr}K$  described above, we derive the asymptotics of  $L$  to be

$$L(r)^{-1} = \mathcal{O}(r^{-2}), \quad \partial_r(L(r)^{-1}) = \mathcal{O}(r^{-3}), \quad |\nabla(L(r)^{-1})| = \mathcal{O}(r^{-3}) \quad (5.46)$$

as  $r$  goes to  $\infty$ .

It follows that  $A$  and  $H$  have continuous first derivatives and satisfy

$$|A(r)| = \mathcal{O}(r^{-3}), \quad |\partial_r A(r)| = \mathcal{O}(r^{-4}), \quad |\nabla A(r)| = \mathcal{O}(r^{-4}) \quad (5.47)$$

$$H(r) = \mathcal{O}(r^{-2}), \quad \partial_r H(r) = \mathcal{O}(r^{-3}), \quad |\nabla H(r)| = \mathcal{O}(r^{-3}) \quad (5.48)$$

By virtue of the fact that  $(g, u, X)$  is a modified solution, we have

$$\partial_r^2 u + \text{tr}K \partial_r u + \mathbb{A}_{g(r)} u = 0 \quad (5.49)$$

The above equation together with the fact that  $u \in \mathcal{A}_\delta^{(2,k+1)}(M)$  then implies  $\partial_r^2 u \in H_{\delta-2}^1([nm_0, \infty); H^{k-2}(S^2))$ . This in particular implies that  $\partial_r u, |\nabla u| \in H_{\delta-1}^2([nm_0, \infty); H^{k-2}(S^2))$ . Using the Sobolev embeddings again and the fact that  $k \geq 5$ , we deduce that  $\partial_r u, |\nabla u|$  have continuous first derivatives and satisfy

$$\partial_r u = o(r^{\delta-1}), \quad \partial_r^2 u = o(r^{\delta-2}), \quad |\nabla \partial_r u| = o(r^{\delta-2}) \quad (5.50)$$

Having achieved the asymptotics for  $u$ , we can now derive an explicit expression for the leading order term for  $H$  and  $\nabla H$ . Using the Gauss equation, we get

$$\begin{aligned} r^2 H(r) &= r^2 Rg(r) - 2 + 2r^2 (\partial_r u)^2 + r^2 |\hat{K}|^2 - \frac{r^2}{2} \left( \text{tr} K - \frac{2}{r} \right)^2 + 2r \left( \text{tr} K - \frac{2}{r} \right) \\ & \quad (5.51) \end{aligned}$$

$$= r^2 R_{g(r)} - 2 + o(r^\delta) \quad (5.52)$$

where we have used equations (5.50) and (5.45).

It then follows that

$$H(r) = r^{-2} (R_{\gamma_\infty} - 2) + o(r^{\delta-2}), \quad |\nabla H(r)| = r^{-3} \left( |R_{\gamma_\infty} - 2|^2 + |\mathcal{R}_{\gamma_\infty}|_{\gamma_\infty}^2 \right)^{1/2} + o(r^{\delta-3}) \quad (5.53)$$

where  $|\cdot|_{\gamma_\infty}$  is the  $\gamma_\infty$ -norm.

In view of the expression of the *Ric* in terms of  $u$ ,  $A$ , and  $H$  in equations (5.34) and (5.35), we deduce the desired regularity of *Ric*, namely that it is  $\mathcal{C}^1$  away from the boundary. We are now in a position to prove equation (5.38). Using again equations (5.34) and (5.35), we estimate  $|Ric|$  and  $|\nabla Ric|$ : for some universal constant  $C > 0$ , we have

$$|Ric| \leq C(|\nabla u|^2 + |A| + |H|), \quad |\nabla Ric| \leq C(|\nabla^2 u|^2 + |\nabla A| + |\nabla H|) \quad (5.54)$$

Using the asymptotics of  $u$ ,  $A$ , and  $H$  laid out in equations (5.50), (5.47) and (5.53), we get that for  $R > nm_0$ ,

$$\sup_{r>R} r^2 (|Ric| + r|\nabla Ric|) \leq C \sup_{S^2} (|R_{\gamma_\infty} - 2| + |\not{d}R_{\gamma_\infty}|_{\gamma_\infty}) + o(R^\delta) \quad (5.55)$$

$$\leq C \|\gamma_\infty - \gamma_{S^2}\|_{\mathcal{C}^3(S^2)} + o(R^\delta) \quad (5.56)$$

$$\leq C \|\gamma_\infty - \gamma_{S^2}\|_{\mathcal{H}^k(S^2)} + o(R^\delta) \quad (5.57)$$

as  $R$  goes to  $\infty$ . In the last line, we have used again the Sobolev embeddings and the fact that  $k \geq 5$ . We have also allowed the constant  $C$  to change from line to line while staying universal, i.e. independent of  $g$  and  $u$ .

We turn our attention to the second statement of the proposition. It suffices to show that  $X_0$  admits 3 derivatives. By virtue of the fact that  $(g, u, X)$  is a modified solution, we have

$$\partial_r \text{tr} K + \frac{1}{2} \text{tr} K^2 + |\hat{K}|^2 = -2(\partial_r u)^2 \quad (5.58)$$

$$\mathcal{L}_{\frac{\partial}{\partial r}} \hat{K} + \text{tr} K \hat{K} = -2 \not{d}u \otimes \not{d}u - g(r)((\partial_r u)^2 - |\nabla u|^2) \quad (5.59)$$

Thanks to equation (5.50), we have that  $|\nabla u| \in H_{\delta-1}^2([nm_0, \infty); H^{k-2}(S^2))$  and so admits 2 radial derivatives. Hence, the above equations directly impy that  $\text{tr} K$  and  $\hat{K}$  admit 3 radial derivative and, in fact, live in  $H_{\delta-1}^3([nm_0, \infty); H^k(S^2))$  and  $H_{\delta-1}^3([nm_0, \infty); \mathcal{H}^k(S^2))$  respectively. Due to the evolution equation

$$\partial_r g(r) = \text{tr} K g(r) + 2\hat{K}, \quad (5.60)$$

we deduce that  $g(r) \in H_{loc}^4([nm_0, \infty); \mathcal{H}^k(S^2))$  and so admits 4 radial derivatives. By the Sobolev embedding and the fact that  $k \geq 5$ , this implies that  $g(r), \partial_r g(r), \partial_r^2 g(r)$  are continuous in  $r$  and are  $\mathcal{C}^3$  on the sphere. We conclude that the metric  $g = dr^2 + g(r)$  is of class  $\mathcal{C}^3$ . Since  $F(X)X$  satisfy the elliptic equation  $\Delta_{g, \text{conf}} F(X)X = 0$  with respect to a  $\mathcal{C}^3$  metric, standard localized interior estimates show that  $F(X)X$  lives in  $\mathcal{X}_\delta^3(M)$  (see for example [CBo8] appendix II).  $\square$

*Remark 5.13.* Stronger regularity results can be proven for the modified solution  $(g, u, X)$ . Specifically,  $g, u,$  and  $X$  are in fact smooth away from the boundary. Nonetheless, the above regularity result is sufficient for our purpose. We will use it to show the nonexistence of nontrivial conformal Killing fields on  $(M, g)$  (see lemma (5.14)).

We now show that  $X = 0$ . Letting  $\bar{X} := F(X)X$ , we note that  $\Delta_{g,conf}\bar{X} = 0$  on  $M$  in light of the fourth line in the definition of  $\Phi$  and the fact that  $(g, u, X)$  is a modified solution. We decompose  $X$  as follows,

$$X = X_0 + X_\infty \quad (5.61)$$

where  $X_0 \in \widehat{\mathcal{X}}_\delta^2(M)$  and  $X_\infty \in \mathcal{X}_\infty$ . We make the following observations:

- If  $\limsup_{r \rightarrow \infty} r^4 |X|^2 = \infty$ , then  $F(X)$  decays exponentially in  $r$ . Since  $|X| = |X_\infty + X_0| = O(r^2)$  by lemma 5.3, we deduce that  $|\bar{X}|$  decays exponentially in  $r$ .
- If  $\limsup_{r \rightarrow \infty} r^4 |X|^2 < \infty$ , then  $F(X) = O(1)$  and so  $|\bar{X}| = O(r^{-2})$ .

The above implies that  $|\bar{X}| = O(r^{-2})$ . This allows us to perform the following integration-by-parts computation:

$$0 = \int_M \bar{X}^\mu \Delta_{g,conf} \bar{X}_\mu dV_g \quad (5.62)$$

$$= -\frac{1}{2} \int_M |\mathcal{L}_{g,conf} \bar{X}|^2 dV_g - \int_{\partial M} \mathcal{L}_{g,conf} \bar{X} \left( \bar{X}, \frac{\partial}{\partial r} \right) d\sigma_{g(nm_0)} \quad (5.63)$$

The above calculation is valid since  $|\mathcal{L}_{g,conf} \bar{X}|^2$  is integrable and the boundary integral at infinity vanishes.

We now compute  $\mathcal{L}_{g,conf} \bar{X} \left( \bar{X}, \frac{\partial}{\partial r} \right)$  on  $\partial M$  in order to evaluate the boundary integral in equation (5.63). Recall that we decomposed  $X$  as follows,

$$X = X_0 + X_\infty \quad (5.64)$$

where  $X_0 \in \widehat{\mathcal{X}}_\delta^2(M)$  and  $X_\infty \in \mathcal{X}_\infty$ . We also have that  $X_0$  and  $X_\infty$  satisfy the following on  $\partial M$ .

$$X_\infty = X_{CK}, \quad \left( \mathcal{L}_{\frac{\partial}{\partial r}} X_0 \right)^T = 0 \quad (5.65)$$

$$\mathcal{L}_{\frac{\partial}{\partial r}} X_\infty = f'(nm_0) \mathit{div}_{\gamma_{S^2}}(X_{CK}) \frac{\partial}{\partial r} + h'(nm_0) X_{CK} = \frac{1}{2} \mathit{div}_{\gamma_{S^2}}(X_{CK}) \frac{\partial}{\partial r} \quad (5.66)$$

$$g\left(X_0, \frac{\partial}{\partial r}\right) = g\left(X_\infty, \frac{\partial}{\partial r}\right) = 0 \quad (5.67)$$

Using the above, we compute that the integrand of the boundary integral appearing in equation (5.63) satisfies the following on  $\partial M$ .



$$\mathcal{L}_{g,conf} \bar{X}(\bar{X}, \frac{\partial}{\partial r}) = \mathcal{L}_{g,conf} X(X, \frac{\partial}{\partial r}) \quad (5.68)$$

$$= \mathcal{L}_X g(X, \frac{\partial}{\partial r}) - \frac{2}{3} \operatorname{div} X g(X, \frac{\partial}{\partial r}) \quad (5.69)$$

$$= g(\mathcal{L}_{\frac{\partial}{\partial r}} X, X) \quad (5.70)$$

$$= g(\mathcal{L}_{\frac{\partial}{\partial r}} X_0, X) + g(\mathcal{L}_{\frac{\partial}{\partial r}} X_\infty, X) \quad (5.71)$$

$$= 0 \quad (5.72)$$

where the fact that  $\bar{X} = X$  near  $\partial M$  was used in the first equality. It then follows that

$$0 = \frac{1}{2} \int_M |\mathcal{L}_{g,conf} \bar{X}|^2 dV_g \quad (5.73)$$

implying that  $\bar{X}$  is conformal Killing on  $(M, g)$ . However, the next lemma shows that if  $\gamma_\infty$  is close enough to  $\gamma_{S^2}$ , then there does not exist a non trivial conformal Killing field on  $(M, g)$  that vanishes at  $\infty$ .

**Lemma 5.14.** *Let  $\delta < 0$ . There exists  $\epsilon > 0$  such that the following holds. Let  $g \in M_\delta^k(M)$  satisfy the statement of proposition (5.12) (i.e. Ric is  $C^1$  away from  $\partial M$  and equation (5.38) holds). Suppose also that  $\|\gamma_\infty - \gamma_{S^2}\|_{\mathcal{H}^k(S^2)} < \epsilon$ . If  $Z \in \mathcal{X}_\delta^3(M)$  is a conformal Killing vector field on  $(M, g)$ , then  $Z = 0$ .*

*Proof.* Let  $Z \in \mathcal{X}_\delta^3(M)$  is a conformal Killing vector field on  $(M, g)$ . A direct computation shows that we can express the third covariant derivative of  $Z$  as follows.

$$\nabla^3 Z = A \cdot \nabla Z + B \cdot Z \quad (5.74)$$

where  $A$  and  $B$  are linear expressions in  $Riem$  and  $\nabla Riem$ , where  $Riem$  is the Riemann curvature tensor. We will move the proof to the appendix to avoid digressing from the main discussion (refer to A.1). Since the dimension of  $M$  is 3,  $Riem$  can be written in terms of only Ric and  $g$ , and so  $A$  and  $B$  can be thought of as linear expressions in  $Ric$  and  $\nabla Ric$ .

An application of a Hardy-type inequality shows that there exists an  $R_0 > nm_0$  depending only on  $g$  and a positive constant  $C$  depending only on  $\delta$  such that for any  $R \geq R_0$  and any vector field  $Z \in \mathcal{X}_\delta^3(M)$ ,

$$\int_{[R,\infty) \times S^2} r^{-2\delta-3} (|Z|^2 + r^2 |\nabla Z|^2) dV_g \leq C \int_{[R,\infty) \times S^2} r^{-2(\delta-3)-3} |\nabla^3 Z|^2 dV_g \quad (5.75)$$

A proof of this inequality is provided in section A.2 in the Appendix (see corollary A.3).

On the other hand, given  $R > nm_0$ , equation (5.74) implies

$$\int_{[R, \infty) \times S^2} r^{-2(\delta-3)-3} |\nabla^3 Z|^2 dV \leq C \int_{[R, \infty) \times S^2} r^{-2(\delta-3)-3} (|\nabla Ric|^2 |Z|^2 + |Ric|^2 |\nabla Z|^2) dV \quad (5.76)$$

$$\leq C \left( \sup_{r > R} r^4 (|Ric|^2 + r^2 |\nabla Ric|^2) \right) \int_{[R, \infty) \times S^2} r^{-2\delta-3} (|Z|^2 + r^2 |\nabla Z|^2) dV \quad (5.77)$$

$$\leq C \left( \|\gamma_\infty - \gamma_{S^2}\|_{\mathcal{H}^k(S^2)}^2 + R^{2\delta} \right) \int_{[R, \infty) \times S^2} r^{-2\delta-3} (|Z|^2 + r^2 |\nabla Z|^2) dV \quad (5.78)$$

$$\leq C(\epsilon + R^{2\delta}) \int_{[R, \infty) \times S^2} r^{-2\delta-3} (|Z|^2 + r^2 |\nabla Z|^2) dV \quad (5.79)$$

where the constant  $C$  is allowed to change from line third to fourth line while staying universal, *i.e.* independent of  $Z$ ,  $R$ , and  $g$ .

We have then proven that

$$\int_{[R, \infty) \times S^2} r^{-2\delta-3} (|Z|^2 + r^2 |\nabla Z|^2) dV \leq C(\epsilon + R^{2\delta}) \int_{[R, \infty) \times S^2} r^{-2\delta-3} (|Z|^2 + r^2 |\nabla Z|^2) dV \quad (5.80)$$

Choosing  $\epsilon$  small enough and  $R$  large enough so that  $C(\epsilon + R^{2\delta}) < 1$  implies that  $Z = 0$  on  $[R, \infty) \times S^2$ .

Since  $Z$  satisfies the elliptic equation  $\Delta_{g, conf} Z = 0$  and vanishes on an open set, standard arguments then imply that  $Z = 0$  on  $M$  (see for example [CO81]).  $\square$

*Remark 5.15.* The nonexistence of nontrivial conformal Killing vector fields vanishing at infinity on asymptotically flat manifolds has already been established in [CO81] and [Yor74]. The above lemma extends this nonexistence result to a broader class of metrics, including some that are not asymptotically flat.

After possibly shrinking the neighbourhood  $\mathcal{U}$  of  $(\gamma_{g_{sc}}, \frac{1}{2} tr K_{g_{sc}})$  and using the continuity of  $\mathcal{H}$ , we can assume that for  $(\gamma_{\mathfrak{B}}, \frac{1}{2} tr K_{\mathfrak{B}}) \in \mathcal{U}$ , the metric  $g$  of the modified solution  $\mathcal{H}(\gamma_{\mathfrak{B}}, \frac{1}{2} tr K_{\mathfrak{B}}) = (g, u, X)$  satisfies

$\|\gamma_\infty - \gamma_{S^2}\|_{\mathcal{H}^k(S^2)} < \epsilon$ . Since  $\bar{X}$  is conformal Killing on  $(M, g)$  and lives in  $\mathcal{X}_\delta^3(M)$ , the above lemma then implies that  $\bar{X} = 0$ , and hence  $X = 0$ . We then finally conclude, by remark 5.7, that  $(g, u)$  is a solution to the conformal static vacuum Einstein equations with Bartnik data  $(\gamma_{\mathfrak{B}}, \frac{1}{2}trK_{\mathfrak{B}})$ . This concludes the proof of the main theorem.

#### 5.4 PROOF THAT $D\Phi_{sc}$ IS AN ISOMORPHISM

In this section, we will prove proposition 5.9.

We first remind the reader of the values of some key parameters for the Schwarzschild solution  $(g_{sc}, f_{sc})$  and the conformal Schwarzschild solution  $g_{sc} = f_{sc}^2 g_{sc}$ ,  $u_{sc} = \ln(f_{sc})$

$$\begin{aligned}
\bullet \quad g_{sc} &= \left(1 - \frac{2m_0}{r}\right)^{-1} dr^2 + r^2 \gamma_{S^2} & \bullet \quad \gamma_{g_{sc}} &= (nm_0)^2 \gamma_{S^2} \\
\bullet \quad trK_{g_{sc}} &= \frac{2\sqrt{1 - \frac{2m_0}{r}}}{r} & \bullet \quad f_{sc} &= \sqrt{1 - \frac{2m_0}{r}} \\
\bullet \quad g_{sc} &= dr^2 + \left(1 - \frac{2m_0}{r}\right) r^2 \gamma_{S^2} & \bullet \quad trK_{sc} &= \frac{2(r-m_0)}{r(r-2m_0)} \\
\bullet \quad \gamma_{sc} &= n(n-2)m_0^2 \gamma_{S^2} & \bullet \quad \hat{K}_{sc} &= 0 \\
\bullet \quad u_{sc} &= \ln \sqrt{1 - \frac{2m_0}{r}} & \bullet \quad R_{\partial M_{sc}} &= \frac{2}{n(n-2)m_0^2}
\end{aligned}$$

Let  $\tilde{g} \in T_{g_{sc}} \mathcal{M}_\delta^k$ ,  $\tilde{u} \in \mathcal{A}_\delta^{(2,k+1)}(M)$ , and  $\tilde{X} \in \hat{\mathcal{X}}_\delta^2(M)$ . For small  $t$ , let  $g(t)$ ,  $u(t)$ , and  $X(t)$  be smooth 1-parameter families satisfying

$$\begin{aligned}
\bullet \quad g(0) &= g_{sc} & \bullet \quad g'(0) &= \tilde{g} \\
\bullet \quad u(0) &= u_{sc} & \bullet \quad u'(0) &= \tilde{u} \\
\bullet \quad X(0) &= 0 & \bullet \quad X'(0) &= \tilde{X}
\end{aligned}$$

Define the following

$$\begin{aligned}
\bullet \quad \widetilde{trK} &:= \left. \frac{d}{dt} \right|_{t=0} trK(t) & \bullet \quad \tilde{\gamma} &:= \left. \frac{d}{dt} \right|_{t=0} \tilde{g}(t)(nm_0) \\
\bullet \quad \tilde{K} &:= \left. \frac{d}{dt} \right|_{t=0} \hat{K}(t) & \bullet \quad \tilde{\omega} &:= \left. \frac{d}{dt} \right|_{t=0} \omega(g(t), X_\infty(t))
\end{aligned}$$

where  $X_\infty(t)$  is the projection of  $X(t)$  into the space  $\mathcal{X}_\infty$ . By definition of  $\omega$ , we have that  $\tilde{\omega}$  is a conformal Killing field on  $(S^2, g_{sc}(nm_0))$ .

We compute  $D\Phi_{sc}$  to be

$$\begin{aligned}
D\Phi_{sc} &: T_{g_{sc}}\mathcal{M}_\delta^k \times \mathcal{A}_\delta^{(2,k+1)}(M) \times \left( \widehat{\mathcal{X}}_\delta^2(M) \oplus \mathcal{X}_\infty(M) \right) \\
&\rightarrow \mathcal{A}_{\delta-2}^{(0,k-1)}(M) \times L_{\delta-2}^2\left([nm_0, \infty); H^k(S^2)\right) \times L_{\delta-2}^2\left([nm_0, \infty); \mathcal{H}^k(S^2)\right) \\
&\quad \times \mathcal{X}_{\delta-2}^0(M) \times H^{k-1}(\partial M) \times \Omega^{k-1}(\partial M) \times \mathcal{H}^k(\partial M) \times H^k(\partial M)
\end{aligned}$$

$$\begin{aligned}
D\Phi_{sc}(\tilde{g}, \tilde{u}, \tilde{X}) &= \frac{d}{dt} \Big|_{t=0} \Phi(\gamma_{g_{sc}}, \frac{1}{2} trK_{g_{sc}}, g(t), u(t), X(t)) \\
&= \begin{pmatrix} \Delta_{g_{sc}} \tilde{u} + (\partial_r u_{sc})(\widetilde{trK}) \\ \partial_r \widetilde{trK} + trK_{sc} \widetilde{trK} + 4(\partial_r u_{sc})(\partial_r \tilde{u}) \\ \mathcal{L}_{\frac{\partial}{\partial r}} \tilde{K} \\ \Delta_{g_{sc}, conf} \tilde{X} \\ -4(\partial_r u_{sc})(\partial_r \tilde{u}) + trK_{sc} \widetilde{trK} \Big|_{\partial M} + \frac{4}{n(n-2)m_0^2} \tilde{u} + 2\mathcal{A}_{\gamma_{sc}} \tilde{u} \\ 2(\partial_r u_{sc}) d\tilde{u} - d\tilde{\sigma}(\tilde{K}) + \tilde{\omega} \\ \frac{n}{n-2} \tilde{\gamma} - 2n^2 m_0^2 \tilde{u} g_{S^2} \\ \widetilde{trK} \Big|_{\partial M} + \frac{2}{nm_0} \tilde{u} - 2\partial_r \tilde{u} \end{pmatrix} \tag{5.81}
\end{aligned}$$

Let  $(A, B, C, D, E, F, G, H)$  be an arbitrary element in the codomain of  $D\Phi_{sc}$ . We wish to show that there exists a unique  $(\tilde{g}, \tilde{u}, \tilde{X})$  in the domain satisfying

$$\Delta_{g_{sc}} \tilde{u} + (\partial_r u_{sc})(\widetilde{trK}) = A, \quad \text{on } M \tag{5.82}$$

$$\partial_r \widetilde{trK} + trK_{sc} \widetilde{trK} + 4(\partial_r u_{sc})(\partial_r \tilde{u}) = B, \quad \text{on } M \tag{5.83}$$

$$\mathcal{L}_{\frac{\partial}{\partial r}} \tilde{K} = C, \quad \text{on } M \tag{5.84}$$

$$\Delta_{g_{sc,conf}} \tilde{X} = D, \quad \text{on } M \quad (5.85)$$

$$-4(\partial_r u_{sc})(\partial_r \tilde{u}) + \text{tr} K_{sc} \widetilde{\text{tr} K} \Big|_{\partial M} + \frac{4}{n(n-2)m_0^2} \tilde{u} + 2\mathbb{A}_{\gamma_{sc}} \tilde{u} = E, \quad \text{on } \partial M \quad (5.86)$$

$$2(\partial_r u_{sc}) \sharp \tilde{u} - \text{div}(\tilde{K}) + \tilde{\omega} = F, \quad \text{on } \partial M \quad (5.87)$$

$$\frac{n}{n-2} \tilde{\gamma} - 2n^2 m_0^2 \tilde{u} g_{S^2} = G, \quad \text{on } \partial M \quad (5.88)$$

$$\widetilde{\text{tr} K} \Big|_{\partial M} + \frac{2}{nm_0} \tilde{u} - 2\partial_r \tilde{u} = H, \quad \text{on } \partial M \quad (5.89)$$

The above equations can be decoupled to give a non-local elliptic system on  $\tilde{u}$ .

**Lemma 5.16.** *Let  $\tilde{u} \in \mathcal{A}_\delta^{(2,k+1)}(M)$  and  $\tilde{g} \in T_{g_{sc}} \mathcal{M}_\delta^k$  satisfy equations (5.83) and (5.89).*

*Then  $\tilde{u}$  and  $\tilde{g}$  satisfy equations (5.82) and (5.86) if and only if  $\tilde{u}$  satisfies*

$$\Delta_{g_{sc}} \tilde{u} - \frac{4m_0^2}{[r(r-2m_0)]^2} \tilde{u} = -\frac{n(n-2)m_0}{2} \frac{4m_0^2}{[r(r-2m_0)]^2} \left( \frac{4-n}{n(n-2)m_0} \tilde{u} \Big|_{\partial M} + \partial_r \tilde{u} \Big|_{\partial M} \right) + \psi, \quad \text{on } M \quad (5.90)$$

$$\frac{4}{nm_0} \partial_r \tilde{u} + 2\mathbb{A} \tilde{u} + \frac{4}{n^2(n-2)m_0^2} \tilde{u} = \Gamma, \quad \text{on } \partial M \quad (5.91)$$

where  $\psi$  and  $\Gamma$  are defined as follows

$$\psi := A - \frac{m_0}{r^2(r-2m_0)^2} \int_{nm_0}^r s(s-2m_0) B ds - \frac{m_0^3 n(n-2)}{r^2(r-2m_0)^2} H \quad (5.92)$$

$$\Gamma := E - \frac{2(n-1)}{n(n-2)m_0} H \quad (5.93)$$

*Remark 5.17.* In light of the spaces that  $A$ ,  $B$ ,  $E$  and  $H$  live in, it follows that  $\psi \in \mathcal{A}_{\delta-2}^{(0,k-1)}(M)$  and  $\Gamma \in H^{k-1}(S^2)$ . Note that the boundary value problem that  $\tilde{u}$  satisfies in equations (5.90) and (5.91) does not depend on  $\tilde{g}$  or  $\tilde{X}$ ; hence, we have indeed decoupled the system.

*Proof.* We directly deduce that equations (5.91) and (5.86) are equivalent by using equation (5.89).

We rewrite equation (5.83) as follows:

$$\partial_r \left( \exp \left( \int_{nm_0}^r \text{tr} K_{sc}(s) ds \right) \widetilde{\text{tr}K} \right) = - \exp \left( \int_{nm_0}^r \text{tr} K_{sc}(s) ds \right) \left( 4(\partial_r u_{sc})(\partial_r \tilde{u}) - B \right). \quad (5.94)$$

A direct computation gives:

$$\exp \left( \int_{nm_0}^r \text{tr} K_{sc}(s) ds \right) = \frac{1}{n(n-2)m_0^2} r(r-2m_0). \quad (5.95)$$

We integrate equation (5.94) to get an expression for  $\widetilde{\text{tr}K}$  in terms of  $\tilde{u}$ : for  $r \in [nm_0, \infty)$  and  $p \in S^2$ , we have

$$\begin{aligned} \widetilde{\text{tr}K}(r, p) &= \frac{n(n-2)m_0^2}{r(r-2m_0)} \widetilde{\text{tr}K} \Big|_{\partial M} (p) - 4 \frac{n(n-2)m_0^2}{r(r-2m_0)} \int_{nm_0}^r \frac{1}{n(n-2)m_0^2} s(s-2m_0) (\partial_r u_{sc})(s) (\partial_r \tilde{u})(s, p) ds \\ &\quad + \frac{n(n-2)m_0^2}{r(r-2m_0)} \int_{nm_0}^r \frac{1}{n(n-2)m_0^2} s(s-2m_0) B(s, p) ds \end{aligned} \quad (5.96)$$

$$\begin{aligned} &= \frac{n(n-2)m_0^2}{r(r-2m_0)} \widetilde{\text{tr}K} \Big|_{\partial M} (p) - 4 \frac{m_0^2}{r(r-2m_0)} \int_{nm_0}^r \frac{1}{m_0^2} s(s-2m_0) \left( \frac{m_0}{s(s-2m_0)} \right) (\partial_r \tilde{u})(s, p) ds \\ &\quad + \frac{1}{r(r-2m_0)} \int_{nm_0}^r s(s-2m_0) B(s, p) ds \end{aligned} \quad (5.97)$$

$$\begin{aligned} &= \frac{n(n-2)m_0^2}{r(r-2m_0)} \widetilde{\text{tr}K} \Big|_{\partial M} (p) - \frac{4m_0}{r(r-2m_0)} \int_{nm_0}^r \partial_r \tilde{u}(s, p) ds + \frac{1}{r(r-2m_0)} \int_{nm_0}^r s(s-2m_0) B(s, p) ds \end{aligned} \quad (5.98)$$

$$\begin{aligned} &= \frac{n(n-2)m_0^2}{r(r-2m_0)} \widetilde{\text{tr}K} \Big|_{\partial M} (p) - \frac{4m_0}{r(r-2m_0)} \left( \tilde{u}(r, p) - \tilde{u}|_{\partial M}(p) \right) \\ &\quad + \frac{1}{r(r-2m_0)} \int_{nm_0}^r s(s-2m_0) B(s, p) ds \end{aligned} \quad (5.99)$$

We plug this into equation (5.82) to derive

$$A = \Delta_{sc}\tilde{u} + \frac{m_0}{r(r-2m_0)} \left( \frac{n(n-2)m_0^2}{r(r-2m_0)} \widehat{trK}|_{\partial M} - \frac{4m_0}{r(r-2m_0)} (\tilde{u} - \tilde{u}|_{\partial M}) \right) + \frac{m_0}{r^2(r-2m_0)^2} \int_{nm_0}^r s(s-2m_0)B(s,p)ds \quad (5.100)$$

$$= \Delta_{sc}\tilde{u} - \frac{4m_0^2}{r^2(r-2m_0)^2} \tilde{u} + \frac{4m_0^2}{r^2(r-2m_0)^2} \left( \tilde{u}|_{\partial M} + \frac{n(n-2)m_0}{4} \widehat{trK}|_{\partial M} \right) + \frac{m_0}{r^2(r-2m_0)^2} \int_{nm_0}^r s(s-2m_0)B(s,p)ds \quad (5.101)$$

Using equation (5.89), it then follows that  $\tilde{u}$  satisfies:

$$\Delta_{g_{sc}}\tilde{u} - \frac{4m_0^2}{r^2(r-2m_0)^2} \tilde{u} = A - \frac{n(n-2)m_0}{2} \frac{4m_0^2}{r^2(r-2m_0)^2} \left( \frac{4-n}{n(n-2)m_0} \tilde{u}|_{\partial M} + \partial_r \tilde{u}|_{\partial M} \right) - \frac{n(n-2)m_0^3}{r^2(r-2m_0)^2} H - \frac{m_0}{r^2(r-2m_0)^2} \int_{nm_0}^r s(s-2m_0)B(s,p)ds \quad (5.102)$$

$$= -\frac{n(n-2)m_0}{2} \frac{4m_0^2}{r^2(r-2m_0)^2} \left( \frac{4-n}{n(n-2)m_0} \tilde{u}|_{\partial M} + \partial_r \tilde{u}|_{\partial M} \right) + \psi \quad (5.103)$$

This proves that equation (5.90) is equivalent to equation (5.82).  $\square$

The rest of the proof will proceed in the following steps.

- Step 1:** We will show that for every  $\psi \in \mathcal{A}_{\delta-2}^{(0,k-1)}(M)$  and  $\Gamma \in H^{k-1}(S^2)$ , there exists a unique solution  $\tilde{u} \in \mathcal{A}_{\delta}^{(2,k+1)}$  solving equations (5.90) and (5.91).
- Step 2:** We will show if  $\tilde{u} \in \mathcal{A}_{\delta}^{(2,k+1)}$  satisfies (5.90) and (5.91) with  $\psi$  and  $\Gamma$  given by equations (5.92) and (5.93), then there exists a unique  $\tilde{g} \in T_{g_{sc}}\mathcal{M}_{\delta}^k(M)$  and a unique conformal Killing field  $\tilde{\omega}$  satisfying equations (5.82) to (5.84) and (5.86) to (5.89).
- Step 3:** We will show that there exists a unique vector field  $\tilde{Y} \in \hat{\mathcal{X}}_{\delta}^2(M)$  satisfying  $\Delta_{g_{sc},conf}\tilde{Y} = D$ .

The above 3 steps will then imply that there exists a unique  $(\tilde{g}, \tilde{u}, \tilde{X})$  in the domain of  $D\Phi_{sc}$  solving equations (5.82) to (5.89). In particular,  $\tilde{u}$  and  $\tilde{g}$  are achieved from steps 1 and 2 respectively and  $\tilde{X} := \tilde{Y} + \tilde{\omega}$ , where  $\tilde{Y}$  and  $\tilde{\omega}$  are achieved from steps 2 and 3 respectively.

### Step 1: Solving for $\tilde{u}$

To study the boundary value problem in (5.90) and (5.91), we will investigate the properties of the corresponding non-local elliptic operator  $\mathcal{P}_{sc}$ , which maps  $\mathcal{A}_\delta^{(2,k+1)}(M)$  into  $\mathcal{A}_{\delta-2}^{(0,k-1)}(M) \times H^{k-1}(\partial M)$  and is defined by

$$\mathcal{P}_{sc}(\tilde{u}) := \begin{pmatrix} \Delta_{g_{sc}} \tilde{u} - \frac{4m_0^2}{[r(r-2m_0)]^2} \tilde{u} + \frac{n(n-2)m_0}{2} \frac{4m_0^2}{[r(r-2m_0)]^2} \left( \frac{4-n}{n(n-2)m_0} \tilde{u}|_{\partial M} + \partial_r \tilde{u}|_{\partial M} \right) \\ \frac{2}{nm_0} \partial_r \tilde{u} + \mathbb{A} \tilde{u} + \frac{2}{n^2(n-2)m_0^2} \tilde{u} \end{pmatrix} \quad (5.104)$$

In fact, this operator will turn out to be Fredholm of index 0 as shown in the following proposition.

*Remark 5.18.* In [AH21] and [AH22b], the authors study the static Einstein vacuum equations in a gauge different from the one used in this paper. Specifically, they study an operator analogous to the operator  $\Phi$  considered here. They achieve that the linearization of their operator is an isomorphism, so as to invoke the implicit function theorem, by first establishing that it is Fredholm of index 0 and then showing that its kernel is trivial. Our approach here is similar except that our gauge allows us to decouple the equations; this decoupling reduces the task of proving that  $D\Phi_{sc}$  is an isomorphism to proving that a much simpler operator,  $\mathcal{P}_{sc}$  acting on the linearization of the lapse function  $\tilde{u}$ , is an isomorphism. Specifically, we will establish that  $\mathcal{P}_{sc}$  is Fredholm of index 0 and has a trivial kernel. The remaining parameters,  $\tilde{g}$  and  $\tilde{X}$ , are governed by straightforward ODEs with  $\tilde{u}$  appearing in the forcing term, and the fact that  $D\Phi_{sc}$  is an isomorphism will follow readily (see **Step 2**).

**Proposition 5.19.** Fix  $\delta \in (-1, -\frac{1}{2}]$  and  $k \in \mathbb{Z}_{\geq 0}$ . Let  $T : \mathcal{A}_\delta^{(2,k+1)}(M) \rightarrow H^{k-1}(\partial M)$  and  $S : \mathcal{A}_\delta^{(2,k+1)}(M) \rightarrow H^{k-1}(\partial M)$  be operators defined by

$$T(\tilde{u}) := \partial_r \tilde{u} + \mu \tilde{u} \quad (5.105)$$

$$S(\tilde{u}) := \mathbb{A}_{\gamma_{sc}} \tilde{u} + \beta_1 \partial_r \tilde{u} + \beta_2 \tilde{u} \quad (5.106)$$

where  $\mu, \beta_1, \beta_2 \in H^k(\partial M)$ .

Let  $\mathcal{P}$  be the nonlocal elliptic differential operator defined by

$$\mathcal{P} : \mathcal{A}_\delta^{(2,k+1)}(M) \rightarrow \mathcal{A}_{\delta-2}^{(0,k-1)}(M) \times H^{k-1}(\partial M)$$



$$\mathcal{P}(\tilde{u}) := \begin{pmatrix} \Delta_{g_{sc}} \tilde{u} - V_1 \tilde{u} - V_2 \overline{T(\tilde{u})} \\ S(\tilde{u}) \end{pmatrix} \quad (5.107)$$

where  $V_1, V_2 \in H_{-3}^1([nm_0, \infty); H^k(S^2))$  and  $\overline{T(\tilde{u})}$  is the function on  $M$  defined by  $(r, p) \mapsto T(\tilde{u})(p)$  for  $(r, p) \in M$ . Then  $\mathcal{P}$  is Fredholm of index 0.

*Proof.* Decompose the operator  $\mathcal{P} = \mathcal{P}_1 + \mathcal{P}_2$ , where the operators  $\mathcal{P}_1, \mathcal{P}_2 : \mathcal{A}_\delta^{(2,k+1)}(M) \rightarrow \mathcal{A}_{\delta-2}^{(0,k-1)}(M) \times H^{k-1}(\partial M)$  are defined by

$$\mathcal{P}_1(\tilde{u}) := \begin{pmatrix} \Delta_{g_{sc}} \tilde{u} \\ \mathbb{A}_{\gamma_{sc}} \tilde{u} \end{pmatrix}, \quad \mathcal{P}_2(\tilde{u}) := \begin{pmatrix} -V_1 \tilde{u} - V_2 \overline{T(\tilde{u})} \\ \beta_1 \partial_r \tilde{u} + \beta_2 \tilde{u} \end{pmatrix}$$

In light of proposition (2.10), we observe that  $\mathcal{P}_2$  is a compact operator. Indeed, for any  $\tilde{u} \in \mathcal{A}_\delta^{(2,k+1)}$ , we have that  $-V_1 \tilde{u} - V_2 \overline{T(\tilde{u})}$  lives in  $H_{-3}^1([nm_0, \infty); H^k(S^2))$  which compactly embeds in  $\mathcal{A}_{\delta-2}^{(0,k-1)}(M)$ , and  $\beta_1 \partial_r \tilde{u} + \beta_2 \tilde{u}$  lives in  $H^k(\partial M)$  which compactly embeds in  $H^{k-1}(\partial M)$ . To show that  $\mathcal{P}$  is Fredholm of index 0, it then suffices to show that  $\mathcal{P}_1$  is Fredholm of index 0 (see [Sch71])

By theorem 3.1, the operator  $\mathcal{Q}$  defined by:

$$\mathcal{Q}(\tilde{u}) := \begin{pmatrix} \Delta_{g_{sc}} \tilde{u} \\ \tilde{u} \end{pmatrix}$$

is an isomorphism from  $\mathcal{A}_\delta^{(2,k+1)}$  to  $\mathcal{A}_{\delta-2}^{(0,k-1)}(M) \times H^{k+1}(\partial M)$ .

We also recall the following standard result on the laplacian on compact manifolds: The operator  $\mathbb{A}_{\gamma_{sc}} : H^{k+1}(\partial M) \rightarrow H^{k-1}(\partial M)$  is Fredholm of index 0. In fact, the kernel is the one-dimensional space of constant functions on  $\partial M$  and the cokernel is the same since  $\mathbb{A}_{\gamma_{sc}}$  is self-adjoint.

We observe that  $\mathcal{P}_1 = (\mathcal{ID}, \mathbb{A}_{\gamma_{sc}}) \circ \mathcal{Q}$ , where

$$(\mathcal{ID}, \mathbb{A}_{\gamma_{sc}}) : \mathcal{A}_{\delta-2}^{(0,k-1)}(M) \times H^{k+1}(\partial M) \rightarrow \mathcal{A}_{\delta-2}^{(0,k-1)}(M) \times H^{k-1}(\partial M)$$

is defined by

$$(\mathcal{ID}, \mathbb{A}_{\gamma_{sc}})(\tilde{v}, f) := (\tilde{v}, \mathbb{A}_{\gamma_{sc}} f)$$

Since  $(\mathcal{ID}, \mathbb{A}_{\gamma_{sc}})$  is Fredholm of index 0 and  $\mathcal{Q}$  is an isomorphism, it then follows that  $\mathcal{P}_1$  is Fredholm of index 0 as needed.  $\square$

By the above proposition, showing that the nonlocal operator  $\mathcal{P}_{sc}$  defined in (5.104) has trivial kernel is sufficient to prove that the system in (5.90) and (5.91) is uniquely solvable for every  $\psi \in \mathcal{A}_{\delta-2}^{(0,k-1)}(M)$  and  $\Gamma \in H^{k-1}(S^2)$ . This will be the content of the next proposition.

**Proposition 5.20.** Fix  $\delta \in (-1, -\frac{1}{2}]$ . Let  $u \in \mathcal{A}_{\delta}^{(2,k+1)}$  satisfy

$$\begin{cases} \Delta_{g_{sc}} \tilde{u} - \frac{4m_0^2}{[r(r-2m_0)]^2} \tilde{u} = -\frac{n(n-2)m_0}{2} \frac{4m_0^2}{[r(r-2m_0)]^2} \left( \frac{4-n}{n(n-2)m_0} \tilde{u}|_{\partial M} + \partial_r \tilde{u}|_{\partial M} \right), & \text{on } M \\ \frac{4}{nm_0} \partial_r \tilde{u} + 2\mathbb{A} \tilde{u} + \frac{4}{n^2(n-2)m_0^2} \tilde{u} = 0, & \text{on } \partial M \end{cases} \quad (5.108)$$

Then  $\tilde{u} = 0$ .

*Proof.* Similarly to what was done in the proof of lemma (3.3), we utilize the spherical symmetry of the conformal Schwarzschild metric to reduce the system in (5.90) and (5.91) to differential equations on the coefficients of  $\tilde{u}$  with respect to its spherical harmonics decomposition:

$$\tilde{u}(r, x) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \tilde{a}_{m\ell}(r) Y_{m\ell}(x) \quad (5.109)$$

for  $r \in [nm_0, \infty)$  and  $x \in \partial M$ .

We define the functions  $a_{m\ell}(r) := \tilde{a}_{m\ell}(nm_0 r)$  on  $[1, \infty)$ . Using the discussion in lemma (3.3), the condition  $\|\tilde{u}\|_{\mathcal{A}_{\delta}^{(2,k+1)}} < \infty$  in particular implies  $\|a_{m\ell}\|_{1,\delta} < \infty$  for every  $m$  and  $\ell$ , where

$$\|a_{m\ell}\|_{1,\delta}^2 = \int_1^{\infty} r^{-2\delta+1} (a'_{m\ell}(r))^2 dr + \int_1^{\infty} r^{-2\delta-1} (a_{m\ell}(r))^2 dr \quad (5.110)$$

The system in (5.90) and (5.91) as well as the condition  $\|a_{m\ell}\|_{1,\delta} < \infty$  implies the following non-local differential equations on all the coefficients  $\tilde{a}_{m\ell}(r)$ :

$$\left\{ \begin{array}{l} r(r-2m_0)\tilde{a}''_{m\ell}(r) + 2(r-m_0)\tilde{a}'_{m\ell}(r) - \left( \frac{4m_0^2}{r(r-2m_0)} + \ell(\ell+1) \right) \tilde{a}_{m\ell}(r) \\ \quad = -\frac{2n(n-2)m_0^3}{r(r-2m_0)} \left( \frac{4-n}{n(n-2)m_0} \tilde{a}_{m\ell}(nm_0) + \tilde{a}'_{m\ell}(nm_0) \right), \quad r \in [nm_0, +\infty) \\ \frac{2}{nm_0} \tilde{a}'_{m\ell}(nm_0) - \frac{1}{n(n-2)m_0^2} (\ell(\ell+1) - \frac{2}{n}) \tilde{a}_{m\ell}(nm_0) = 0, \\ \|a_{m\ell}\|_{1,\delta} < \infty \end{array} \right. \quad (5.111)$$

The functions  $a_{m\ell}$ , using (5.111), satisfy the following similar non-local differential equations:

$$\left\{ \begin{array}{l} r(r-\frac{2}{n})a''_{m\ell}(r) + 2(r-\frac{1}{n})a'_{m\ell}(r) - \left( \frac{4}{n^2r(r-\frac{2}{n})} + \ell(\ell+1) \right) a_{m\ell}(r) \\ \quad = -\frac{2(n-2)}{n^2r(r-\frac{2}{n})} \left( \frac{4-n}{n-2} a_{m\ell}(1) + a'_{m\ell}(1) \right), \\ \frac{2}{n^2} a'_{m\ell}(1) - \frac{1}{n(n-2)} (\ell(\ell+1) - \frac{2}{n}) a_{m\ell}(1) = 0, \\ \|a_{m\ell}\|_{1,\delta} < \infty \end{array} \right. \quad (5.112)$$

Note that the mass parameter  $m_0$  does not appear in the non-local differential equation for  $a_{m\ell}$ . From here onwards we will study the system (5.112) instead of (5.111).

We consider (5.112) for any nonnegative integer  $\ell$  and seek to derive that  $a_{m\ell}(r) = 0$  is the only solution. We consider the  $\ell = 0$  case separately. We replace the last condition in (5.112) with  $a'_{m\ell}(1) = C$  and find the explicit (unique) solution to this shooting problem to be

$$a_{00}(r) = -C \frac{-2+n+6r-3nr-2nr^2+n^2r^2}{r(nr-2)}, \quad (5.113)$$

It then can easily be verified that  $\lim_{r \rightarrow \infty} a_{00}(r) = 0$  if and only if  $C = 0$ , implying that  $a_{00} = 0$  is the only solution to (5.112).

We conclude the only solutions to (5.112) in the  $\ell = 0$  case is the zero solution.

We now deal with the  $\ell \geq 1$  case. We will write  $a_\ell$  instead of  $a_{m\ell}$  for simplicity.

Define the following constants:

$$\bullet \quad C_\ell = -\frac{2(n-2)}{n^2} \left( \frac{4-n}{n-2} a_\ell(1) + a'_\ell(1) \right)$$

$$\bullet \quad \alpha_\ell = \frac{n^2}{n(2-\ell(\ell+1))-6}$$

$$\bullet \quad \beta_\ell = \frac{n^2(n\ell(\ell+1)-2)}{(n(2-\ell(\ell+1))-6)(2(n-2))}$$

Note that  $\alpha_\ell$  and  $\beta_\ell$  are well defined for  $\ell \geq 1$  and are both negative. Then the function  $a_\ell$  solves the following initial value problem:

$$\begin{cases} r(r - \frac{2}{n})a''_\ell(r) + 2(r - \frac{1}{n})a'_\ell(r) - \left( \frac{4}{n^2 r(r - \frac{2}{n})} + \ell(\ell+1) \right) a_\ell(r) = \frac{1}{r(r - \frac{2}{n})} C_\ell \\ a'_\ell(1) = \beta_\ell C_\ell \\ a_\ell(1) = \alpha_\ell C_\ell \end{cases} \quad (5.114)$$

If  $C_\ell = 0$ , then  $a_\ell = 0$  by the existence and uniqueness theorem from ODE theory. Suppose now that  $C_\ell \neq 0$ . By considering  $\frac{a_\ell}{C_\ell}$  instead of  $a_\ell$ , we can assume without loss of generality that  $C_\ell = 1$ .

We rewrite the system in the following way:

$$\begin{cases} \frac{d}{dr} \left[ r(r - \frac{2}{n})a'_\ell(r) \right] = \left( \frac{4}{n^2 r(r - \frac{2}{n})} + \ell(\ell+1) \right) a_\ell(r) + \frac{1}{r(r - \frac{2}{n})} \\ a'_\ell(1) = \beta_\ell \\ a_\ell(1) = \alpha_\ell \end{cases} \quad (5.115)$$

We will show that  $a_\ell$  blows up at infinity contradicting that  $\|a_\ell\|_{1,\delta} < \infty$ . This will then imply that  $C_\ell = 0$  and hence  $a_\ell = 0$ . First, we prove a technical lemma.

**Lemma 5.21.** *Let  $h_1, h_2$  be smooth functions on  $[1, \infty)$  such that  $h_1$  is positive and  $\lim_{r \rightarrow \infty} h_1(r) = C_1$  for some  $C_1 > 0$ . Let  $g$  be a function on  $[1, \infty)$  satisfying the following ODE:*

$$\frac{d}{dr} \left[ r(r - \frac{2}{n})g'(r) \right] = h_1(r)g(r) + h_2(r) \quad (5.116)$$

Then the following is true.

- Suppose that  $h_2(r) \geq 0$  ( $\leq 0$ ) on  $[1, \infty)$  and that both  $g(r_*)$  and  $g'(r_*)$  are positive (negative) for some  $r_* \in [1, \infty)$ . Then  $g$  and  $g'$  are positive (negative) on  $(r_*, \infty)$ .
- Suppose that  $g$  and  $g'$  are positive (negative) on  $(r_*, \infty)$  for some  $r_* \in [1, \infty)$  and that  $h_2(r) = O(r^{-2})$ . Then  $\lim_{r \rightarrow \infty} g(r) = \infty$  ( $-\infty$ ).

*Proof.* Suppose that  $h_2$  is nonnegative everywhere. Let  $r_* \in [1, \infty)$  be such that  $g'(r_*) > 0$  and  $g(r_*) > 0$ . We will prove that  $g'(r) > 0$  on  $(r_*, \infty)$  by using a simple bootstrap method. By continuity of  $g'(r)$ , we know that  $g'(r) > 0$  on  $[r_*, r_* + \delta)$  for some  $\delta > 0$ . Then the set  $B := \{r \in (r_*, \infty) \mid g'(s) > 0 \text{ for } s \in [r_*, r)\}$  is nonempty. Suppose that  $R := \sup B < \infty$ . By continuity, we have that  $g'(R) = 0$  and  $g'(r) > 0$  for  $r \in (r_*, R)$ . Since  $g(r_*) > 0$  and  $g$  is increasing on  $(r_*, R)$ , we have that  $g(R) > 0$ . By letting  $r = R$  in equation (5.116), it follows that  $\frac{d}{dr}[r(r - 2/n)g']$  is positive at  $R$  and, in turn, on a neighbourhood of  $R$ . This implies that  $r(r - 2/n)g'(r)$  is increasing on a neighbourhood of  $R$ , which implies that  $0 < r(r - 2/n)g'(r) < R(R - 2/n)g'(R)$  for  $r < R$  and close to  $R$ . As this contradicts that  $g'(R) = 0$ , we conclude that  $\sup B = \infty$  and hence  $g'$  and  $g$  are positive on  $(r_*, \infty)$ .

Suppose now that  $g$  and  $g'$  are positive on  $(r_*, \infty)$  for some  $r_* \in [1, \infty)$  and that  $h_2(r) = O(r^{-2})$ . In virtue of the positivity of  $g'$  and  $g$  as well as the monotone convergence theorem, it follows that  $\lim_{r \rightarrow \infty} g(r)$  either is a positive number or is  $\infty$ . Suppose that  $\lim_{r \rightarrow \infty} g(r) = A > 0$ , which in particular implies that  $g'$  is integrable.

By integrating equation (5.116), we get that

$$r(r - 2/n)g'(r) = \frac{n-2}{n}g'(1) + \int_1^r [h_1(s)g(s) + h_2(s)]ds \quad (5.117)$$

Using the fact that  $h_1$  and  $g$  are  $O(1)$  and that  $h_2(r) = O(r^{-2})$ , it follows that  $\sup_{r \geq 1} r g'(r) < \infty$ .

Furthermore, we have that

$$rg'(r) - g'(1) = \int_1^r [sg'(s)]' ds \quad (5.118)$$

$$= \int_1^r g'(s) ds + \int_1^r sg''(s) ds \quad (5.119)$$

$$= \int_1^r \frac{h_1(s)g(s)}{s - 2/n} ds + \int_1^r \frac{h_2(s)}{s - 2/n} ds - \int_1^r \frac{s}{s - 2/n} g'(s) ds \quad (5.120)$$

It follows that there exists a constant  $M > 0$  such that

$$\int_1^r \frac{h_1(s)g(s)}{s - 2/n} ds \leq M \quad (5.121)$$

for any  $r \in [1, \infty)$ ; since the integrand is positive, the limit as  $r$  tends to  $\infty$  exists. In particular, this implies that  $\frac{g}{r} \in L^1([1, \infty))$ , which then contradicts that  $A$  is positive. Hence, we see that  $\lim_{r \rightarrow \infty} g(r) = \infty$ , as needed.

The case when  $h_2$  is nonpositive is identical. □

**Corollary 5.22.** *Let  $h_1, h_2$  be smooth functions on  $[1, \infty)$  such that  $h_1$  is positive and  $\lim_{r \rightarrow \infty} h_1(r) = C_1$  for some  $C_1 > 0$ . Let  $g$  and  $\tilde{g}$  be functions on  $[1, \infty)$  satisfying the following ODE:*

$$\frac{d}{dr} \left[ r \left( r - \frac{2}{n} \right) g'(r) \right] = h_1(r)g(r) + h_2(r), \quad \frac{d}{dr} \left[ r \left( r - \frac{2}{n} \right) \tilde{g}'(r) \right] = h_1(r)\tilde{g}(r) + h_2(r), \quad (5.122)$$

If  $g(1) = \tilde{g}(1)$  and  $g'(1) < \tilde{g}'(1)$ , then  $g(r) < \tilde{g}(r)$  for any  $r \in (1, \infty)$ .

*Proof.* Define  $f := g - \tilde{g}$  and observe that  $f$  satisfies

$$\frac{d}{dr} \left[ r \left( r - \frac{2}{n} \right) f'(r) \right] = h_1(r)f(r) \quad (5.123)$$

Observe also that  $f(1) = 0$  and  $f'(1) < 0$ . This in particular implies that there exists an  $r_* > 1$  close enough to 1 such that  $f(r)$  and  $f'(r)$  are negative for any  $r \in (1, r_*)$ . By invoking lemma (5.21), we conclude that  $f(r) < 0$  for any  $r \in (1, \infty)$  as needed. □

We now return to our goal of showing that  $a_\ell$  blows up at  $\infty$ .

We first decompose  $a_\ell = f_\ell + g_\ell$  where  $f_\ell$  solves

$$\begin{cases} \frac{d}{dr} [r(r - \frac{2}{n})f'_\ell(r)] = \left(\frac{4}{n^2r(r - \frac{2}{n})} + \ell(\ell + 1)\right) f_\ell(r) - \frac{4\alpha_\ell}{n^2r(r - 2/n)} \\ f'_\ell(1) = 0 \\ f_\ell(1) = \alpha_\ell \end{cases} \quad (5.124)$$

and  $g_\ell$  solves

$$\begin{cases} \frac{d}{dr} [r(r - \frac{2}{n})g'_\ell(r)] = \left(\frac{4}{n^2r(r - \frac{2}{n})} + \ell(\ell + 1)\right) g_\ell(r) + \frac{1 + (4/n^2)\alpha_\ell}{r(r - 2/n)} \\ g'_\ell(1) = \beta_\ell \\ g_\ell(0) = 0 \end{cases} \quad (5.125)$$

By letting  $r = 1$  in the equation for  $f_\ell$ , we observe that  $r(r - 2/n)f'_\ell(r)$  is decreasing near  $r = 1$ ; since  $f'_\ell(1) = 0$ , it follows that  $f_\ell$  and  $f'_\ell$  are negative near  $r = 1$ . In particular, letting  $\tilde{f}_\ell(r) := f_\ell(r) - \alpha_\ell$ , we also have that  $\tilde{f}_\ell$  and  $\tilde{f}'_\ell$  are negative near  $r = 1$ . Using the system in (5.124), we deduce that  $\tilde{f}_\ell$  satisfies the following:

$$\begin{cases} \frac{d}{dr} [r(r - \frac{2}{n})\tilde{f}'_\ell(r)] = \left(\frac{4}{n^2r(r - \frac{2}{n})} + \ell(\ell + 1)\right) \tilde{f}_\ell(r) + \alpha_\ell\ell(\ell + 1) \\ \tilde{f}'_\ell(1) = 0 \\ \tilde{f}_\ell(1) = 0 \end{cases} \quad (5.126)$$

We invoke lemma (5.21) on  $\tilde{f}_\ell$  to deduce that  $\tilde{f}_\ell$  and  $\tilde{f}'_\ell$  are negative on  $(1, \infty)$ . This in particular implies that  $f_\ell(r)$  and  $f'_\ell(r)$  are negative on  $(1, \infty)$ . We again invoke lemma (5.21) on  $f_\ell$  to conclude that  $\lim_{r \rightarrow \infty} f_\ell(r) = -\infty$ .

It suffices to show that  $g_\ell(r) < 0$  for all  $r \in [1, \infty)$ . We first observe from the definition of  $\beta_\ell$  and  $\alpha_\ell$  that

$$\frac{\beta_\ell}{1 + (4/n^2)\alpha_\ell} = -\frac{n^2}{2(n-2)} \left(1 + \frac{2(n-2)}{n(\ell(\ell+1) - 2) + 2}\right) \quad (5.127)$$

$$< -\frac{n^2}{2(n-2)} \quad (5.128)$$

for all  $n > 2$  and  $\ell \in \mathbb{N}$ . In particular, we have that  $1 + (4/n^2)\alpha_\ell > 0$  for every  $n > 2$  and  $\ell \in \mathbb{N}$ .

Let  $\tilde{g}_\ell$  be the function on  $[1, \infty)$  solving

$$\begin{cases} \frac{d}{dr} \left[ r \left( r - \frac{2}{n} \right) \tilde{g}'_\ell(r) \right] = \left( \frac{4}{n^2 r \left( r - \frac{2}{n} \right)} + \ell(\ell + 1) \right) \tilde{g}_\ell(r) + \frac{1}{r \left( r - \frac{2}{n} \right)} \\ \tilde{g}'_\ell(1) = -\frac{n^2}{2(n-2)} \\ \tilde{g}_\ell(1) = 0 \end{cases} \quad (5.129)$$

In light of corollary (5.22) along with equation (5.128), it follows that the negativity of  $\tilde{g}_\ell(r)$  on  $(1, \infty)$  implies the negativity of  $\frac{g_\ell(r)}{1 + (4/n^2)\alpha_\ell}$  on  $(1, \infty)$ , which in turn implies that the negativity of  $g_\ell(r)$  on  $(1, \infty)$ . It then suffices to show that  $\tilde{g}_\ell(r) < 0$  on  $(1, \infty)$ .

We will prove that  $\tilde{g}_\ell(r) < 0$  on  $(1, \infty)$  by induction on  $\ell$ . We first find the solution for  $\ell = 0$  to be:

$$\tilde{g}_0(r) = -\frac{n(r-1)}{2(r-2/n)} \quad (5.130)$$

which is negative everywhere.

Now suppose that  $\tilde{g}_\ell(r) < 0$  on  $(1, \infty)$  for some nonnegative integer  $\ell$ . Define  $h_\ell := \tilde{g}_{\ell+1} - \tilde{g}_\ell$ , which will solve:

$$\begin{cases} \frac{d}{dr} \left[ r \left( r - \frac{2}{n} \right) h'_\ell(r) \right] = \left( \frac{4}{n^2 r \left( r - \frac{2}{n} \right)} + (\ell + 1)(\ell + 2) \right) h_\ell(r) + 2(\ell + 1)\tilde{g}_\ell(r) \\ h'_\ell(1) = 0 \\ h_\ell(1) = 0 \end{cases} \quad (5.131)$$

We directly compute  $h''_\ell(1) = \frac{n}{n-2} 2(\ell + 1)\tilde{g}_\ell(1) = 0$  and  $h'''_\ell(1) = \frac{n}{n-2} 2(\ell + 1)\tilde{g}'_\ell(1) < 0$ . This then implies that  $h_\ell$  and  $h'_\ell$  are negative near  $r = 1$ . Using the fact that  $\tilde{g}_\ell(r) < 0$  on  $(1, \infty)$  and invoking lemma (5.21), it follows that  $h_\ell(r) < 0$  on  $(1, \infty)$ , which in turn implies that  $\tilde{g}_{\ell+1}(r) < 0$  on  $(1, \infty)$  as needed.

We have finally shown that  $a_\ell$  blows up at infinity for every  $\ell \in \mathbb{N}$  contradicting that  $\|a_\ell\|_{1,\delta} < \infty$ . We conclude that the assumption that  $C_\ell \neq 0$  was false and hence  $a_\ell = 0$  for every  $\ell \in \mathbb{N}$  and, in turn,  $\tilde{u} = 0$ .  $\square$

This concludes Step 1, and we move on to Step 2.



## Step 2: Solving for $\tilde{g}$ and $\tilde{\omega}$

Let  $\tilde{u} \in \mathcal{A}_\delta^{(2,k+1)}$  be the function satisfying equations (5.90) and (5.91) with  $\psi$  and  $\Gamma$  given by equations (5.92) and (5.93). We wish to show that there exists a unique  $\tilde{g} \in T_{g_{sc}}\mathcal{M}_\delta^k$  and a conformal Killing field  $\tilde{\omega}$  satisfying equations (5.82) to (5.84) and (5.86) to (5.89).

Equation (5.89) determines uniquely the initial data for  $\widetilde{trK}$ , which is given by

$$\widetilde{trK}\Big|_{\partial M} = H + 2\partial_r\tilde{u}\Big|_{\partial M} - \frac{2}{nm_0}\tilde{u}\Big|_{\partial M} \quad (5.132)$$

and is living in  $H^k(\partial M)$ . We rewrite the ODE (5.83) obeyed by  $\widetilde{trK}$  here for convenience.

$$\partial_r\widetilde{trK} + trK_{sc}\widetilde{trK} + 4(\partial_r u_{sc})(\partial_r\tilde{u}) = B, \quad \text{on } M$$

which, together with the initial condition in (5.132), determines uniquely  $\widetilde{trK}$  on  $M$ . We explicitly solve for  $\widetilde{trK}$  on  $M$  to get:

$$\widetilde{trK}(r, p) = \frac{n(n-2)m_0^2}{r(r-2m_0)} \widetilde{trK}\Big|_{\partial M}(p) - \frac{4m_0}{r(r-2m_0)} (\tilde{u}(r, p) - \tilde{u}\Big|_{\partial M}(p)) + \frac{1}{r(r-2m_0)} \int_{nm_0}^r s(s-2m_0)B(s, p)ds \quad (5.133)$$

for  $r \in [nm_0, \infty)$  and  $p \in S^2$ .

We observe that

- $(\partial_r u_{sc})(\partial_r\tilde{u}) \in L_{\delta-2}^2([nm_0, \infty); H^k(S^2))$
- $B \in L_{\delta-2}^2([nm_0, \infty); H^k(S^2))$ .

and so  $\widetilde{trK}$  lies in  $H_{\delta-1}^1([nm_0, \infty); H^k(S^2))$ .

We turn our attention to  $\tilde{K}$ . We first recall the well known fact regarding the divergence operator on symmetric traceless tensors on  $S^2$  (see [CK93]).

**Proposition 5.23.** *Let  $k \geq 2$ . Let  $\gamma$  be a smooth metric on  $S^2$  with positive curvature. Denote by  $\mathcal{D}^k(S^2)$  the space of traceless symmetric  $(0, 2)$  tensors on  $(S^2, \gamma)$  with components in  $H^k(S^2)$ . Let  $\Omega^{\perp k-1}(\partial M)$  be the space of vector fields on  $S^2$  with components in  $H^{k-1}(M)$  that are  $L^2$  orthogonal to conformal Killing vector fields on  $(S^2, \gamma)$ . Then the divergence operator  $\text{div}_\gamma$  is an isomorphism from  $\mathcal{D}^k(S^2)$  to  $\Omega^{\perp k-1}(S^2)$*

The above proposition along with equation (5.87) imply that  $\frac{2}{n(n-2)m_0}d\tilde{u} + \tilde{\omega} - F$  must be orthogonal to conformal Killing vector fields on  $(\partial M, \gamma_{sc})$ . Since  $\tilde{\omega}$  is conformal Killing on  $(\partial M, \gamma_{sc})$ , this requirement determines  $\tilde{\omega}$  uniquely. Indeed, if  $Y_1, \dots, Y_6$  is an  $L^2$  orthonormal basis of conformal Killing vector fields, then  $\tilde{\omega}$  must be

$$\tilde{\omega} := \sum_{i=1}^6 \left( \int_{S^2} Y_i \cdot \left( F - \frac{2}{n(n-2)m_0} d\tilde{u} \right) d\sigma_{\gamma_{sc}} \right) Y_i \quad (5.134)$$

The above proposition together with (5.87) determine uniquely the initial condition for  $\tilde{K}$  to be

$$\tilde{K}|_{\partial M} = d\tilde{v}_{\gamma_{sc}}^{-1} \left( \frac{2}{n(n-2)m_0} d\tilde{u} + \tilde{\omega} - F \right) \quad (5.135)$$

living in  $\mathcal{H}^k(\partial M)$ . We rewrite the ODE (5.84) obeyed by  $\tilde{K}$  here for convenience.

$$\mathcal{L}_{\frac{\partial}{\partial r}} \tilde{K} = C, \quad \text{on } M$$

which, together with the initial condition in (5.135), determines  $\hat{K}$  uniquely on  $M$ . Fixing fermi coordinates,  $(r, \theta^1, \theta^2)$ , we explicitly solve for  $\hat{K}$  on  $M$  to get:

$$\hat{K}_{ij}(r, p) = \tilde{K}_{ij}|_{\partial M} + \int_{nm_0}^r C_{ij}(s, p) \quad (5.136)$$

for  $i, j = 1, 2, r \in [nm_0, \infty)$  and  $p \in S^2$ .

Since  $C \in L_{\delta-2}^2([nm_0, \infty); \mathcal{H}^k(S^2))$ , it follows that  $\hat{K}$  lies in  $H_{\delta-1}^1([nm_0, \infty); \mathcal{H}^k(S^2))$ .

Equation (5.88) determines uniquely initial data for  $\tilde{g}$  given by

$$\tilde{g}|_{\partial M} = \frac{n-2}{n} (2n^2 m_0^2 \tilde{u} \gamma_{S^2} + G) \quad (5.137)$$

living in  $\mathcal{H}^k(\partial M)$ . The evolution of  $\tilde{g}$  is determined by  $\widetilde{trK}$  and  $\tilde{K}$  in the following equation:

$$\mathcal{L}_{\frac{\partial}{\partial r}} \tilde{g} = 2\tilde{K} + \widetilde{trK} g_{sc} + trK_{sc} \tilde{g} \quad (5.138)$$

Equations (5.138) and (5.137) determine uniquely  $\tilde{g}$  to be, in fermi coordinates  $(r, \theta^1, \theta^2)$ ,

$$\tilde{g}_{ij} = r(r - 2m_0) \int_{nm_0}^r \frac{1}{s(s - 2m_0)} \left( 2\tilde{K}_{ij} + \widetilde{trK} g_{scij} \right) ds + \frac{r(r - 2m_0)}{nm_0^2} G_{ij} \quad (5.139)$$

In light of the fact that  $\widetilde{trK} \in H_{\delta-1}^1([nm_0, \infty); H^k(S^2))$  and  $\tilde{K} \in H_{\delta-1}^1([nm_0, \infty); \mathcal{H}^k(S^2))$ , it follows  $\tilde{g}$  is of the form  $\tilde{g} = r^2(G + h(r))$  where  $G \in \mathcal{H}^k(S^2)$  and  $h \in H_{\delta}^2([nm_0, \infty); \mathcal{H}^k(S^2))$ ; this implies that  $\tilde{g} \in T_{g_{sc}} \mathcal{M}_{\delta}^k$  as needed.

We have then shown that there exists a unique  $\tilde{g}$  and  $\tilde{\omega}$  satisfying equations (5.83), (5.84) and (5.86) to (5.89). It follows by lemma (5.16) that equation (5.82) is satisfied as well. This concludes Step 2, and we move on to Step 3.

### Step 3: Solving for $\tilde{X}$

We wish to show that there exists a unique  $\tilde{Y} \in \hat{\mathcal{X}}_{\delta}^2(M)$  satisfying

$$\Delta_{g_{sc}, conf} \tilde{Y} = D \quad (5.140)$$

where  $D \in \mathcal{X}_{\delta-2}^0(M)$ . Similar results have been shown in [Max05] for the above equation with trivial Dirichlet and Neumann conditions. In our case, vector fields  $Y$  in  $\hat{\mathcal{X}}^2(M)$  satisfy the following mixed boundary conditions

$$g\left(Y, \frac{\partial}{\partial r}\right) = 0, \quad \left(\mathcal{L}_{\frac{\partial}{\partial r}} Y\right)^T = 0 \quad (5.141)$$

The isomorphism of the operator  $\Delta_{g_{sc}, conf}$  in our space follows by minor modifications of the proof in [Max05]. We add the proof here for the sake of completeness.

**Proposition 5.24.** *Let  $\delta \in (-1, -\frac{1}{2}]$ . The operator  $\Delta_{g_{sc}, conf}$  is an isomorphism from  $\hat{\mathcal{X}}_{\delta}^2(M)$  to  $\mathcal{X}_{\delta-2}^0(M)$ .*

*Proof.* Recall that if  $X \in \hat{\mathcal{X}}_{\delta}^2(M)$ , then

$$g\left(X, \frac{\partial}{\partial r}\right) = 0, \quad \left(\mathcal{L}_{\frac{\partial}{\partial r}} X\right)^T = 0 \quad (5.142)$$

on  $\partial M$ . In particular, we have that

$$\mathcal{L}_{g_{sc},conf}X(X, \frac{\partial}{\partial r}) = \mathcal{L}_X g(X, \frac{\partial}{\partial r}) - \frac{2}{3} \operatorname{div} X g(X, \frac{\partial}{\partial r}) \quad (5.143)$$

$$= g(\mathcal{L}_{\frac{\partial}{\partial r}} X, X) \quad (5.144)$$

$$= 0 \quad (5.145)$$

on  $\partial M$ .

Let  $X \in \widehat{\mathcal{X}}_\delta^2(M)$  satisfy  $\Delta_{g_{sc},conf}X = 0$ . Given  $R \geq nm_0$ , let  $\phi_R$  be a cutoff function on  $[nm_0, \infty)$  satisfying  $\phi_R(x) = 1$  for  $x \leq R$ ,  $\phi_R(x) = 0$  for  $x \geq R + 1$ , and  $-2 \leq \phi'_R(x) \leq 0$  for any  $x \in [nm_0, \infty)$ .

We integrate by parts to get

$$0 = \int_M \phi_R X^\mu \Delta_{g_{sc},conf} X_\mu dV_{g_{sc}} \quad (5.146)$$

$$= -\frac{1}{2} \int_M (\mathcal{L}_{g_{sc},conf} \phi_R X) \cdot (\mathcal{L}_{g_{sc},conf} X) dV_{g_{sc}} - \int_{\partial M} \mathcal{L}_{g_{sc},conf} X(X, \frac{\partial}{\partial r}) d\sigma_{g_{sc}(nm_0)} \quad (5.147)$$

$$= -\frac{1}{2} \int_M \phi_R |\mathcal{L}_{g_{sc},conf} X|^2 dV_{g_{sc}} - \frac{1}{2} \int_M 2\phi'_R \mathcal{L}_{g_{sc},conf} X(X, \frac{\partial}{\partial r}) dV_{sc} \quad (5.148)$$

This implies that for any  $R \geq nm_0$ ,

$$\int_{B_R \setminus B_{nm_0}} |\mathcal{L}_{g_{sc},conf} X|^2 dV_{g_{sc}} \leq \int_M \phi_R |\mathcal{L}_{g_{sc},conf} X|^2 dV_{g_{sc}} \quad (5.149)$$

$$= - \int_M 2\phi'_R \mathcal{L}_{g_{sc},conf} X(X, \frac{\partial}{\partial r}) dV_{sc} \quad (5.150)$$

$$\leq 4 \int_{B_{R+1} \setminus B_R} |\mathcal{L}_{g_{sc},conf} X(X, \frac{\partial}{\partial r})| dV_{sc} \quad (5.151)$$

$$\lesssim \int_{B_{R+1} \setminus B_R} |\nabla X|^2 dV_{sc} \quad (5.152)$$

Since  $\delta \in (-1, -\frac{1}{2})$ , we have that  $|\nabla X|^2 \leq |\nabla X|^2 r^{-2\delta-1}$  and so  $|\nabla X|^2$  is integrable on  $M$  and  $\int_M |\nabla X|^2 dV_{sc} \leq \|\nabla X\|_{0,\delta-1}^2$ . We can then take the limit as  $R$  goes to infinity in the above equations to deduce that

$$\int_M |\mathcal{L}_{g_{sc},conf} X|^2 dV_{g_{sc}} = 0 \quad (5.153)$$

implying that  $X$  is conformal Killing on  $(M, g_{sc})$ . Since the equation  $\Delta_{g_{sc},conf}X = 0$  is an elliptic PDE with smooth coefficients, elliptic reg-

ularity shows that  $X$  is  $C^\infty$ . We can then invoke lemma 5.14 to conclude that  $X = 0$ , which shows that the kernel of  $\Delta_{g_{sc},conf}$  is trivial.

Now we show that the kernel of the adjoint is also trivial. It will then follow that  $\Delta_{g_{sc},conf}$  is an isomorphism. Recall that

$$\Delta_{g_{sc},conf}^* : (\mathcal{X}_{\delta-2}^0(M))^* \rightarrow (\mathcal{X}_\delta^2(M))^*$$

For any number  $\tau \in \mathbb{R}$ , Riesz's representation theorem allows us to identify  $(\mathcal{X}_\tau^0(M))^*$  with  $\mathcal{X}_\tau^0(M)$  via the map :

$$J : \mathcal{X}_\tau^0(M) \rightarrow (\mathcal{X}_\tau^0(M))^*$$

$$\text{For } Y \in \mathcal{X}_\tau^0(M), J(Y) : X \in \mathcal{X}_\tau^0(M) \mapsto \int_M X \cdot Y r^{-2\tau-3} dV_{g_{sc}}$$

where  $\cdot$  is with respect to  $g_{sc}$ . For simplicity of the notation, we will denote both  $Y$  and  $J(Y)$  by  $Y$ ; it will be clear from context which one we are referring to.

We then have that  $Y \in (\mathcal{X}_{\delta-2}^0(M))^*$  is in the kernel of  $\Delta_{g_{sc},conf}^*$  if and only if

$$\int_M Y \cdot \Delta_{g_{sc},conf} X r^{-2(\delta-2)-3} dV_{g_{sc}} = 0 \quad (5.154)$$

for every  $X \in \widehat{\mathcal{X}}_\delta^2(M)$ , which is equivalent to the above equation holding for every smooth compactly supported vector field  $X$  in  $\widehat{\mathcal{X}}_\delta^2(M)$  by a density argument.

It follows from elliptic regularity that  $Y \in \mathcal{X}_{\delta-2}^2(M)$ . In fact,  $Y$  will be smooth since the metric  $g_{sc}$  is smooth. Given an arbitrary smooth compactly supported vector field  $X$  in  $\widehat{\mathcal{X}}_\delta^2(M)$ , we can then integrate by parts to get

$$\int_M Y \cdot \Delta_{g_{sc}, conf} X r^{-2(\delta-2)-3} dV_{sc} = \int_M X \cdot \Delta_{g_{sc}, conf} \tilde{Y} dV_{sc} \quad (5.155)$$

$$+ \int_{\partial M} \left( \mathcal{L}_{g_{sc}, conf} \tilde{Y} \left( X, \frac{\partial}{\partial r} \right) - \mathcal{L}_{g_{sc}, conf} X \left( \tilde{Y}, \frac{\partial}{\partial r} \right) \right) d\sigma \quad (5.156)$$

$$= \int_M X \cdot \Delta_{g_{sc}, conf} \tilde{Y} dV_{sc} \quad (5.157)$$

$$\int_{\partial M} \left[ X^i \left( \mathcal{L}_{\frac{\partial}{\partial r}} \tilde{Y}_i - \partial_i \tilde{Y}_0 \right) + \frac{5}{3} \partial_r X_0 \tilde{Y}_0 + \frac{2}{3} \text{div}(X^T) \tilde{Y}_0 \right] d\sigma \quad (5.158)$$

where  $\tilde{Y} := r^{-2(\delta-2)-3} Y$ ,  $X_0 := g_{sc}(\frac{\partial}{\partial r}, X)$  and  $\tilde{Y}_0 := g_{sc}(\frac{\partial}{\partial r}, \tilde{Y})$ . Note that the boundary terms vanish at infinity since  $X$  is compactly supported. Since  $X$  was an arbitrary vector field in a dense subset of  $\hat{\mathcal{X}}_\delta^2(M)$ , it follows that  $\tilde{Y}$  satisfies

$$\Delta_{g_{sc}, conf} \tilde{Y} = 0, \quad g\left(\frac{\partial}{\partial r}, \tilde{Y}\right) = 0, \quad \left(\mathcal{L}_{\frac{\partial}{\partial r}} \tilde{Y}\right)^T = 0 \quad (5.159)$$

Considering that  $Y \in \mathcal{X}_{\delta-2}^0(M)$ , it follows that  $\tilde{Y} \in \hat{\mathcal{X}}_{-\delta-1}^2(M)$ . Since  $\Delta_{g_{sc}, conf} \tilde{Y} = 0$  and  $-1 < -\delta - 1 < 0$ , we have that  $Y \in \hat{\mathcal{X}}_\tau^2(M)$  for any  $\tau \in (-1, 0)$  (see [CBo8] and [Bar86]). We can then apply the same integration-by-parts argument carried out earlier to conclude that  $\tilde{Y}$  is conformal Killing on  $(M, g_{sc})$  and hence, by lemma (5.14), vanishes as needed. □

# APPENDIX

---

## A.1 A PDE OF FINITE TYPE FOR CONFORMAL KILLING VECTOR FIELDS

In this section, we will prove an identity satisfied by conformal Killing vector fields that is used in the proof of lemma 5.14. More specifically, we will prove that any conformal Killing vector field  $Z$  on an arbitrary  $n$ -dimensional Riemannian manifold  $(M, g)$  satisfies the following PDE of finite type:

$$\nabla^3 Z = A \cdot \nabla Z + B \cdot Z \quad (\text{A.1})$$

where  $A$  and  $B$  are linear expressions in  $Riem$  and  $\nabla Riem$ .

Let  $Z$  be a conformal Killing field and let  $\psi := \frac{2}{n} \text{div} Z$ . The conformal Killing equation is

$$\nabla_i Z_j + \nabla_j Z_i = \psi g_{ij} \quad (\text{A.2})$$

Eisenhart in [Eis49] (see pages 231-232 in [Eis49]) proves the following identities:

$$\nabla_k \nabla_j Z_i = -Z_m R_{kij}^m + \frac{1}{2} (g_{ij} \nabla_k \psi + g_{ik} \nabla_j \psi - g_{jk} \nabla_i \psi) \quad (\text{A.3})$$

$$g^{il} Z_m \nabla_l R_{kij}^m - Z_m \nabla_k R_j^m - \nabla_k Z_m R_j^m - \nabla_j Z_m R_k^m + \frac{n-2}{2} \nabla_k \nabla_j \psi + \frac{1}{2} g_{jk} \Delta \psi = 0 \quad (\text{A.4})$$

where  $R$  denotes  $Riem$  or  $Ric$  depending on the number of indices. Taking the trace of equation (A.4), we get

$$\Delta \psi = \frac{2}{n-1} \left( Z_m \nabla_i R^{mi} + \nabla_i Z_m R^{mi} \right) \quad (\text{A.5})$$

Using equation (A.5) to eliminate  $\Delta\psi$  in equation (A.4), we get the following expression of  $\nabla_k\nabla_j\psi$ :

$$\nabla_k\nabla_j\psi = \frac{2}{n-2} \left( -g^{il}Z_m\nabla_l R_{kij}^m + Z_m\nabla_k R_j^m + \nabla_k Z_m R_j^m + \nabla_j Z_m R_k^m - \frac{1}{n-1}g_{jk} \left( Z_m\nabla_i R^{mi} + \nabla_i Z_m R^{mi} \right) \right) \quad (\text{A.6})$$

Taking a derivative of equation (A.3) and using equation (A.6), we get the desired result.

## A.2 A HARDY-TYPE INEQUALITY

In this section, we will prove a Hardy-type inequality that is used in the proof of lemma 5.14, namely equation (5.75).

Let  $g \in \mathcal{M}_\delta^k(M)$  be a Riemannian metric on  $M = [nm_0, \infty) \times S^2$  of the form

$$g = dr^2 + r^2(\gamma_\infty + h(r))$$

where  $\gamma_\infty$  is a metric on  $S^2$  and  $h \in H_\delta^2([nm_0, \infty); \mathcal{H}^k(S^2))$ . We will prove the following.

**Proposition A.1.** *There exists an  $R_0 > nm_0$  depending only on  $g$  such that for  $R \geq R_0$  and  $\tau > 0$ , the inequality*

$$\int_{[R, \infty) \times S^2} r^{\tau-2} |T|^2 dV_g \leq \frac{4}{\tau^2} \int_{[R, \infty) \times S^2} r^\tau |\nabla T|^2 dV_g \quad (\text{A.7})$$

holds for all tensor fields  $T \in C_c^1(M)$ .

*Proof.* The main tool we will use is a general  $L^p$  Hardy inequality in Riemannian manifolds developed by D'Ambrosio and Dipierro in [DD12]. We present the relevant version of it below.

**Theorem A.2.** *Let  $\rho \in C^2(M)$  such that  $\Delta_g \rho \geq 0$  such that  $\frac{|\nabla \rho|^2}{\Delta \rho} \in L_{loc}^1(M)$ . Then for any  $R > nm_0$ , the inequality*

$$\int_{[R, \infty) \times S^2} |u|^2 \Delta \rho dV_g \leq 4 \int_{[R, \infty) \times S^2} \frac{|\nabla \rho|^2}{\Delta \rho} |\nabla u|^2 dV_g \quad (\text{A.8})$$

holds for all  $u \in C_c^1(M)$ .

Letting  $\rho = r^\tau$  for  $\tau > 0$ , we compute

$$\Delta \rho = \tau r^{\tau-2} \left[ \tau + 1 + r \left( \text{tr}K - \frac{2}{r} \right) \right], \quad |\nabla \rho|^2 = \tau^2 r^{2\tau-2} \quad (\text{A.9})$$



In light of the Sobolev embeddings in proposition 2.10, we have that

$$\left| \operatorname{tr}K - \frac{2}{r} \right| = o(r^{-1+\delta}) \quad (\text{A.10})$$

So we can choose  $R_0 > nm_0$  depending only on  $g$  such that for any  $R \geq R_0$ ,

$$\left| r \left( \operatorname{tr}K - \frac{2}{r} \right) \right| \leq 1 \quad (\text{A.11})$$

This, in turn, implies that for any  $\tau > 0$  and any  $R \geq R_0$ ,

$$\Delta\rho \geq \tau^2 r^{\tau-2} \geq 0 \quad (\text{A.12})$$

on  $[R, \infty) \times S^2$ .

We can then invoke theorem A.2 directly to deduce that the inequality

$$\int_{[R, \infty) \times S^2} r^{\tau-2} |u|^2 dV_g \leq \frac{4}{\tau^2} \int_{[R, \infty) \times S^2} r^\tau |\nabla u|^2 dV_g \quad (\text{A.13})$$

hold for any  $R \geq R_0$  and  $u \in C_c^1(M)$ .

The same inequality holds with  $u$  replaced with a tensor field  $T$ . To see this, we first compute that for any tensor field  $T$  on  $M$ ,

$$|\nabla|T||^2 \leq |\nabla T|^2 \quad (\text{A.14})$$

Using the above and letting  $u = |T|$  in equation (A.13), we immediately deduce that the inequality

$$\int_{[R, \infty) \times S^2} r^{\tau-2} |T|^2 dV_g \leq \frac{4}{\tau^2} \int_{[R, \infty) \times S^2} r^\tau |\nabla T|^2 dV_g \quad (\text{A.15})$$

holds for any  $R \geq R_0$  and tensor field  $T \in C_c^1(M)$ .  $\square$

**Corollary A.3.** *Let  $\delta \in (-1, -\frac{1}{2})$ . There exists an  $R_0 > nm_0$  depending only on  $g$  and a positive constant  $C$  depending only on  $\delta$  such that for any  $R \geq R_0$  and any vector field  $Z \in \mathcal{X}_\delta^3(M)$ ,*

$$\int_{[R, \infty) \times S^2} r^{-2\delta-3} (|Z|^2 + r^2 |\nabla Z|^2) dV_g \leq C \int_{[R, \infty) \times S^2} r^{-2(\delta-3)-3} |\nabla^3 Z|^2 dV_g \quad (\text{A.16})$$

*Proof.* By repeatedly applying proposition A.1 for  $\tau = -2\delta - 1, -2\delta + 1, -2\delta + 3$  and  $T = Z, \nabla Z, \nabla^2 Z$ , we deduce that there exists a positive constant  $C = C(\delta)$  such that

$$\int_{[R,\infty) \times S^2} r^{-2\delta-3} (|Z|^2 + r^2 |\nabla Z|^2) dV_g \leq C \int_{[R,\infty) \times S^2} r^{-2(\delta-3)-3} |\nabla^3 Z|^2 dV_g \quad (\text{A.17})$$

for any  $R \geq R_0$  and any vector field  $Z \in C_c^3(M)$ . The above inequality can be rewritten in terms of the norm on  $\mathcal{X}_\delta^k(M)$  (see definition 2.5) as follows:

$$\|Z\|_{1,\delta}^2 \leq C \|\nabla^3 Z\|_{0,\delta-3}^2 \quad (\text{A.18})$$

The desired inequality then follows from the density of  $C_c^3(M)$  in  $\mathcal{X}_\delta^3(M)$ .  $\square$

### A.3 THE LEGENDRE FUNCTIONS $P_\ell$ AND $Q_\ell$

In this section, we will prove properties of the Legendre functions of the first and second kind,  $P_\ell$  and  $Q_\ell$ , that are used in the proof of lemma 3.3.

Fix  $R > 1$  (which is  $n - 1$  in chapter 3). For a positive integer  $\ell$ , as described by Olver in [OR14], the Legendre functions,  $P_\ell$  and  $Q_\ell$ , are linearly independent solutions to the ODE

$$(z^2 - 1)h''(z) + 2zh'(z) - \ell(\ell + 1)h(z) = 0, \quad z \in [R, \infty) \quad (\text{A.19})$$

with the following asymptotics as  $z \rightarrow \infty$ ,

$$P_\ell(z) = O(z^\ell), \quad Q_\ell(z) = O(z^{-\ell-1}) \quad (\text{A.20})$$

We normalize  $P_\ell$  and  $Q_\ell$  so that

$$\lim_{z \rightarrow \infty} z^{-\ell} P_\ell(z) = 1, \quad \lim_{z \rightarrow \infty} z^{\ell+1} Q_\ell(z) = 1 \quad (\text{A.21})$$

which is different than Olver's. Letting  $\mathbf{P}_\ell$  and  $\mathbf{Q}_\ell$  be the Legendre functions as defined by Olver, the relation between ours and his can immediately be obtained is as follows (see [OR14] chapter 5 section 12 and 13):

$$P_\ell(z) = \frac{\sqrt{\pi}\Gamma(\ell + 1)}{2^\ell\Gamma(\ell + \frac{1}{2})} \mathbf{P}_\ell(z), \quad Q_\ell(z) = \frac{2^{\ell+1}\Gamma(\ell + \frac{3}{2})}{\sqrt{\pi}\Gamma(\ell + 1)} \mathbf{Q}_\ell(z) \quad (\text{A.22})$$

In the following proposition, we will apply the method of Frobenius to obtain the expansion of  $P_\ell$  and  $Q_\ell$  in terms of powers of  $z$ .

**Proposition A.4.**  $P_\ell$  and  $Q_\ell$  admit an expansion of the following form. For  $z > 1$ ,

$$P_\ell(z) = \sum_{k=0}^{\ell} a_k z^{\ell-k}, \quad Q_\ell(z) = \sum_{k=0}^{\infty} b_k z^{-\ell-1-k} \quad (\text{A.23})$$

where the coefficients  $a_k$  and  $b_k$  are defined recursively as follows:

$$a_0 = b_0 = 1, \quad a_1 = b_1 = 0$$

$$\text{for } k \geq 2, \quad a_k = \frac{(\ell - k + 2)(\ell - k + 1)}{k^2 - k(2\ell + 1)} a_{k-2}, \quad b_k = \frac{(\ell + k - 1)(\ell + k)}{k(2\ell + k + 1)} b_{k-2}$$

The expansion of  $P_\ell$  and  $Q_\ell$  as described above agree with [WW96] pg 302 and 320.

**Proposition A.5.** There exists a constant  $C = C(R)$  such that for any  $\ell \geq 1$  and  $z \in [R, \infty)$ , the following holds

$$z^{-\ell} |P_\ell(z)| \leq C \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^{-\ell}, \quad z^{\ell+1} |Q_\ell(z)| \leq C \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^{\ell} \quad (\text{A.24})$$

$$z^{-(\ell-1)} |P'_\ell(z)| \leq C\ell \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^{-\ell}, \quad z^{\ell+2} |Q'_\ell(z)| \leq C\ell \left( \frac{2z}{z + \sqrt{z^2 - 1}} \right)^{\ell} \quad (\text{A.25})$$

*Proof.* Olver, in [OR14] chapter 12 section 12, has established an asymptotic expansion of  $\mathbf{P}_\ell$  and  $\mathbf{Q}_\ell$  for large degree  $\ell$  that is uniformly valid for  $z \in (1, \infty)$ . Shivakumar and Wong, in [SW88], proved an equivalent expansion of  $\mathbf{P}_\ell$  that is more computable; letting  $u = \ell + 1/2$ , he has shown that for  $\xi > 0$ ,

$$\mathbf{P}_\ell(\cosh \xi) = \left( \frac{\xi}{\sinh \xi} \right)^{1/2} \left( I_0(u\xi) + \epsilon(u, \xi) \right) \quad (\text{A.26})$$

where

$$|\epsilon(u, \xi)| \leq \frac{\Gamma(3/2)}{2\Gamma(1/2)} \frac{2\xi}{1 + \xi} \frac{I_1(u\xi)}{u} \quad (\text{A.27})$$

and  $I_0, I_1$  are the modified Bessel functions (see [OR14] chapter 2 section 10).

Similarly, Frenzen, in [Fre90], proved an equivalent expansion of  $\mathbf{Q}_\ell$  that is more computable than Olver's; he has shown that for  $\xi > 0$ ,

$$\mathbf{Q}_\ell(\cosh \xi) = \left( \frac{\xi}{\sinh \xi} \right)^{1/2} \left( K_0(u\xi) + \eta(u, \xi) \right) \quad (\text{A.28})$$

where

$$|\eta(u, \zeta)| \leq \frac{\zeta}{2(2 + \zeta)} \frac{K_1(u\zeta)}{u} \quad (\text{A.29})$$

and  $K_0, K_1$  are the modified Bessel functions (see [OR14] chapter 7 section 8)

Letting  $\zeta_0$  be the positive number in which  $\cosh \zeta_0 = R$ , Olver's asymptotics of the modified Bessel function in [OR14] chapter 12, section 1 implies that for all  $\zeta \geq \zeta_0$ ,

$$|I_0(u\zeta)| + |I_{-1}(u\zeta)| \leq C \frac{e^{u\zeta}}{2\pi\sqrt{u\zeta}}, \quad |K_0(u\zeta)| + |K_1(u\zeta)| \leq C \left( \frac{\pi}{2u\zeta} \right)^{1/2} e^{-u\zeta} \quad (\text{A.30})$$

for some constant  $C > 0$  depending only on  $R$ . We can then compute for  $\zeta \in [\zeta_0, \infty)$ ,

$$(\cosh \zeta)^{-\ell} |P_\ell(\cosh \zeta)| = \frac{\sqrt{\pi}\Gamma(\ell + 1)}{2^\ell \Gamma(\ell + \frac{1}{2})} (\cosh \zeta)^{-\ell} |\mathbf{P}_\ell(\cosh \zeta)| \quad (\text{A.31})$$

$$\leq C \frac{\sqrt{\pi}\Gamma(\ell + 1)}{2^\ell \Gamma(\ell + \frac{1}{2})} 2^\ell (e^\zeta + e^{-\zeta})^{-\ell} \frac{\sqrt{2\zeta}}{\sqrt{e^\zeta - e^{-\zeta}}} \frac{e^{(\ell+1/2)\zeta}}{2\pi\sqrt{(\ell+1/2)}\sqrt{\zeta}} \quad (\text{A.32})$$

$$\leq C \frac{\Gamma(\ell + 1)}{\Gamma(\ell + \frac{1}{2})} \frac{1}{\sqrt{\ell}} \left( \frac{e^\zeta}{e^\zeta - e^{-\zeta}} \right)^{1/2} \left( \frac{e^\zeta}{e^\zeta + e^{-\zeta}} \right)^\ell \quad (\text{A.33})$$

$$\leq C \frac{\Gamma(\ell + 1)}{\Gamma(\ell + \frac{1}{2})} \frac{1}{\sqrt{\ell}} \left( \frac{e^\zeta}{e^\zeta + e^{-\zeta}} \right)^\ell \quad (\text{A.34})$$

$$(\cosh \zeta)^{\ell+1} |Q_\ell(\cosh \zeta)| = \frac{2^{\ell+1}\Gamma(\ell + \frac{3}{2})}{\sqrt{\pi}\Gamma(\ell + 1)} (\cosh \zeta)^{\ell+1} |\mathbf{Q}_\ell(\cosh \zeta)| \quad (\text{A.35})$$

$$\leq C \frac{2^{\ell+1}\Gamma(\ell + \frac{3}{2})}{\sqrt{\pi}\Gamma(\ell + 1)} 2^{-\ell-1} (e^\zeta + e^{-\zeta})^{\ell+1} \frac{\sqrt{2\zeta}}{\sqrt{e^\zeta - e^{-\zeta}}} \left( \frac{\pi}{2(\ell+1/2)\zeta} \right)^{1/2} e^{-(\ell+1/2)\zeta} \quad (\text{A.36})$$

$$\leq C \frac{\Gamma(\ell + \frac{3}{2})}{\Gamma(\ell + 1)} \frac{1}{\sqrt{\ell}} \left( \frac{e^\zeta}{e^\zeta - e^{-\zeta}} \right)^{1/2} \left( \frac{e^\zeta + e^{-\zeta}}{e^\zeta} \right)^{\ell+1} \quad (\text{A.37})$$

$$\leq C \frac{\Gamma(\ell + \frac{3}{2})}{\Gamma(\ell + 1)} \frac{1}{\sqrt{\ell}} \left( \frac{e^\zeta + e^{-\zeta}}{e^\zeta} \right)^\ell \quad (\text{A.38})$$

where we have allowed the constant  $C$  to change from line to line but remains dependent only on  $R$  and not on  $\zeta$  or  $\ell$ .

In light of Stirling's formula (see [OR14], chapter 3, section 8), the Gamma function enjoys the following asymptotics:

$$\Gamma(x) = e^{-x}x^x(1 + \mathcal{O}(x^{-1})), \quad \text{as } x \rightarrow \infty \quad (\text{A.39})$$

In particular, we have

$$\frac{\Gamma(\ell+1)}{\Gamma(\ell+\frac{1}{2})} = \left(1 + \frac{1}{2(\ell+1/2)}\right)^{\ell+1/2} \frac{\sqrt{\ell+1}}{\sqrt{e}}(1 + \mathcal{O}(\ell^{-1})) \quad (\text{A.40})$$

and

$$\frac{\Gamma(\ell+\frac{3}{2})}{\Gamma(\ell+1)} = \left(1 + \frac{1}{2(\ell+1)}\right)^{\ell+1} \frac{\sqrt{\ell+\frac{3}{2}}}{\sqrt{e}}(1 + \mathcal{O}(\ell^{-1})) \quad (\text{A.41})$$

as  $\ell \rightarrow \infty$ . It then follows that  $\frac{\Gamma(\ell+1)}{\Gamma(\ell+\frac{1}{2})} \frac{1}{\sqrt{\ell}}$  and  $\frac{\Gamma(\ell+\frac{3}{2})}{\Gamma(\ell+1)} \frac{1}{\sqrt{\ell}}$  are bounded for  $\ell \geq 1$ , and we finally conclude that there exists a constant  $C$  depending only on  $R$  such that for any  $z \geq R$ ,

$$z^{-\ell}P_\ell(z) \leq C \left(\frac{2z}{z + \sqrt{z^2 - 1}}\right)^{-\ell}, \quad z^{\ell+1}Q_\ell(z) \leq C \left(\frac{2z}{z + \sqrt{z^2 - 1}}\right)^\ell \quad (\text{A.42})$$

where we used

$$\frac{e^\xi + e^{-\xi}}{e^\xi} = \frac{2z}{z + \sqrt{z^2 - 1}} \quad (\text{A.43})$$

for  $z = \cosh \xi$ .

We have then shown equation (A.24). The estimate for  $P'_\ell$  and  $Q'_\ell$  in equation (A.25) follows immediately from equation (A.24) and the recurrence relations (see [BE55] pg 161 and [WW96] pg 318)

$$(z^2 - 1)P'_\ell(z) = \ell(zP_\ell(z) - P_{\ell-1}(z)), \quad (z^2 - 1)Q'_\ell(z) = \ell(zQ_\ell(z) - Q_{\ell-1}(z)) \quad (\text{A.44})$$

□

# BIBLIOGRAPHY

---

- [AH21] Zhongshan An and Lan-Hsuan Huang, *Existence of static vacuum extensions with prescribed bartnik boundary data*, Cambridge Journal of Mathematics (2021).
- [AH22a] ———, *New asymptotically flat static vacuum metrics with near euclidean boundary data*, Journal of Mathematical Physics (2022).
- [AH22b] ———, *Static vacuum extensions with prescribed bartnik boundary data near a general static vacuum metric*, Annals of PDE (2022).
- [AK09] Michael T. Anderson and Marcus A. Khuri, *On the bartnik extension problem for the static vacuum einstein equations*, Classical and Quantum Gravity **30** (2009).
- [AMR12] Ralph Abraham, Jerrold E Marsden, and Tudor Ratiu, *Manifolds, tensor analysis, and applications*, vol. 75, Springer Science & Business Media, 2012.
- [And15] Michael T. Anderson, *Local existence and uniqueness for exterior static vacuum einstein metrics*, Proceedings of the American Mathematical Society **143** (2015), no. 7, 3091–3096.
- [And23] ———, *The bartnik quasi-local mass conjectures*, 2023.
- [Bar86] Robert Bartnik, *The mass of an asymptotically flat manifold*, Communications on Pure and Applied Mathematics **39** (1986), 661–693.
- [Bar89] ———, *New definition of quasilocal mass*, Phys. Rev. Lett. **62** (1989), 2346–2348.
- [Bar97] ———, *Energy in general relativity*, Tsing Hua Lectures on Geometry and Analysis (Hsinchu, 1990–1991) (1997), 5—27.
- [BE55] Harry Bateman and Arthur Erdélyi, *Higher transcendental functions*, California Institute of technology. Bateman Manuscript project, McGraw-Hill, New York, NY, 1955.
- [BO13] Carl M Bender and Steven A Orszag, *Advanced mathematical methods for scientists and engineers i: Asymptotic methods and perturbation theory*, Springer Science & Business Media, 2013.

- [CB08] Yvonne Choquet-Bruhat, *General relativity and the einstein equations*, OUP Oxford, 2008.
- [CK93] Demetrios Christodoulou and Sergiu Klainerman, *The global nonlinear stability of the minkowski space*, Séminaire Equations aux dérivées partielles (Polytechnique) (1993), 1–29.
- [CO81] Demetrios Christodoulou and Niall O’Murchadha, *The boost problem in general relativity*, Communications in Mathematical Physics **80** (1981), 271–300.
- [Cor00] J. Corvino, *Scalar curvature deformation and a gluing construction for the einstein constraint equations.*, Comm. Math. Phys. **214** (2000), 137–189.
- [Cor17] ———, *A note on the bartnik mass*, Nonlinear analysis in geometry and applied mathematics, Harv. Univ. Cent. Math. Sci. Appl. Ser. Math., **1**, Int. Press, Somerville, MA. (2017), 45–79.
- [DD12] Lorenzo D’Ambrosio and Serena Dipierro, *Hardy inequalities on riemannian manifolds and applications*, arXiv: Analysis of PDEs (2012).
- [Eis49] L.P. Eisenhart, *Riemannian geometry*, Princeton University Press, 1949.
- [Eva98] L.C. Evans, *Partial differential equations*, Graduate studies in mathematics, American Mathematical Society, 1998.
- [Fre90] C. L. Frenzen, *Error bounds for a uniform asymptotic expansion of the legendre function  $q_n^{-m}(\cosh z)$* , SIAM Journal on Mathematical Analysis **21** (1990), no. 2, 523–535.
- [GT13] D. Gilbarg and N.S. Trudinger, *Elliptic partial differential equations of second order*, Grundlehren der mathematischen Wissenschaften, Springer Berlin Heidelberg, 2013.
- [Isr67] Werner Israel, *Event horizons in static vacuum space-times*, Phys. Rev. **164** (1967), 1776–1779.
- [KM95] Daniel Kennefick and Niall Ó Murchadha, *Weakly decaying asymptotically flat static and stationary solutions to the einstein equations*, Classical and Quantum Gravity **12** (1995), no. 1, 149–158.
- [KN12] Sergiu Klainerman and Francesco Nicolò, *The evolution problem in general relativity*, vol. 25, Springer Science & Business Media, 2012.

- [KW75] Jerry Kazdan and Frank W. Warner, *Existence and conformal deformation of metrics with prescribed gaussian and scalar curvatures*, *Annals of Mathematics* **101** (1975), 317.
- [Max05] David Maxwell, *Solutions of the einstein constraint equations with apparent horizon boundaries*, *Communications in mathematical physics* **253** (2005), no. 3, 561–583.
- [Mia03] Pengzi Miao, *On existence of static metric extensions in general relativity*, *Communications in mathematical physics* **241** (2003), no. 1, 27–46.
- [OR14] F.W.J. Olver and W. Rheinbolt, *Asymptotics and special functions*, Elsevier Science, 2014.
- [Rico2] D. Richards, *Advanced mathematical methods with maple*, *Advanced Mathematical Methods with Maple*, Cambridge University Press, 2002.
- [Sch71] M. Schechter, *Principles of functional analysis*, Student edition, Academic Press, 1971.
- [SW88] P. N. Shivakumar and R. Wong, *Error bounds for a uniform asymptotic expansion of the legendre function  $p_n^{-m}(\cosh z)$* , *Quarterly of Applied Mathematics* **46** (1988), no. 3, 473–488.
- [WW96] E. T. Whittaker and G. N. Watson, *A course of modern analysis*, 4 ed., *Cambridge Mathematical Library*, Cambridge University Press, 1996.
- [Yor74] James W. Jr. York, *Covariant decompositions of symmetric tensors in the theory of gravitation*, *Annales De L Institut Henri Poincare-physique Theorique* **21** (1974), 319–332.